

From the Department of Microbiology, Tumor and Cell Biology  
Karolinska Institutet, Stockholm, Sweden

# **GLYCAN-BASED INTERACTIONS OF STREPTOCOCCUS PNEUMONIAE AND THE HOST**

Karina Hentrich



**Karolinska  
Institutet**

Stockholm 2017

Cover: Fluorescent microscopy picture of A549 lung epithelial cells and *Streptococcus pneumoniae* TIGR4. Sialic acids on the cell surfaces are stained with FITC-labelled *Sambucus nigra* lectin (green), nuclei are stained with DAPI (blue) and pneumococci are labelled with Nile red (red).

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Eprint AB 2017

© Karina Hentrich, 2017

ISBN 978-91-7676-657-6

# GLYCAN-BASED INTERACTIONS OF STREPTOCOCCUS PNEUMONIAE AND THE HOST

AKADEMISK AVHANDLING

som för avläggande av medicine doktorsexamen vid Karolinska Institutet  
offentligen försvaras i **Petrén, Nobels väg 12B**, Karolinska Institutet,  
Solna

**Fredagen den 9 juni 2017, kl. 09.00**

av

**Karina Hentrich**

*Huvudhandledare:*

Professor Birgitta Henriques-Normark  
Karolinska Institutet  
Institutionen för Mikrobiologi, Tumör- och  
Cellbiologi

*Bihandledare:*

Professor Staffan Normark  
Karolinska Institutet  
Institutionen för Mikrobiologi, Tumör- och  
Cellbiologi

Ph.D. Jonas Löfling

AstraZeneca

Global Operations, CMC-RC and Stability

*Fakultetsopponent:*

Professor Marco Rinaldo Oggioni  
University of Leicester  
Department of Genetics

*Betygsnämnd:*

Docent Teresa Frisan  
Karolinska Institutet  
Institutionen för Cell- och Molekylärbiologi

Professor Jan-Ingmar Flock  
Karolinska Institutet  
Institutionen för Mikrobiologi, Tumör- och  
Cellbiologi

Docent Constantin Urban  
Umeå Universitet  
Institutionen för Klinisk Mikrobiologi,  
Immunologi



*Für Moritz*



## ABSTRACT

*Streptococcus pneumoniae* is commonly found as an asymptomatic colonizer of the nasopharynx of children, but it can also translocate to normally sterile body sites and cause severe diseases, like pneumonia, septicemia or meningitis. Pneumococci spread via aerosols. Upon entry into the upper respiratory tract of the host, glycoconjugates with terminal sialic acids (Sias) are among the first structures pneumococci encounter. Hence, they play an important role in pneumococcal pathogenesis. Moreover, glycans are also implicated in the recognition of microbial pathogens by the innate immune system, as many ligands of Toll-like receptors are glycoconjugates. Both aspects of glycan-based pneumococcal-host interactions were studied in this thesis.

In most mammals, the Sia *N*-acetylneuraminic acid (Neu5Ac) is converted into *N*-glycolylneuraminic acid (Neu5Gc) by the cytidine-monophosphate-*N*-acetylneuraminic acid hydroxylase (CMAH). However, humans lack Neu5Gc due to a deletion in *CMAH*, instead they overproduce Neu5Ac. We reported a faster disease progression in *Cmah*<sup>-/-</sup> versus wild-type (wt) mice after pneumococcal challenge and an upregulation of pneumococcal sialidase NanA and the main sialic acid transporter SatABC in response to Neu5Ac as compared with Neu5Gc, which was mediated by the response regulator CiaR.

Moreover, we detected higher pneumococcal adhesion rates to cells presenting Neu5Ac than Neu5Gc. *In vitro*, higher bacterial adherence downregulated IL-8 secretion, and *in vivo*, pneumococcal pyruvate oxidase (SpxB) and pneumolysin contributed to a reduced immune response in *Cmah*<sup>-/-</sup> compared with wt mice after intranasal challenge.

Influenza infections lead to changes in the pulmonary environment and sensitize for a pneumococcal infection. We observed higher protein concentrations, increased numbers of dead cells as well as upregulated hydrogen peroxide concentrations in bronchoalveolar lavages of influenza- versus mock-infected mice. The increased virus-mediated stress in the lower respiratory tract mediated an upregulation of the pneumococcal serine protease HtrA during influenza/pneumococcal coinfection. A mutant of HtrA was severely attenuated in a murine coinfection model, suggesting an important role of HtrA in pneumococcal outgrowth following primary influenza infection.

Dendritic cells link the innate with the adaptive immune system. We found an RNA-mediated recognition of pneumococci by TLR3 in dendritic cells, which induced the secretion of the cytokine IL-12. Moreover, in influenza/pneumococcal coinfections, the virus upregulated TLR3 expression, which led to an enhanced production of IL-12 by dendritic cells.

In summary, we show that glycan-mediated interactions of *S. pneumoniae* and the host play a major role in pneumococcal host tropism and strongly affect pneumococcal virulence, as well as innate immune responses.

## LIST OF SCIENTIFIC PAPERS

This thesis is based on the following publications, which are referred to in the text by Roman numerals:

- I. **HENTRICH K.**, LÖFLING J., PATHAK A., NIZET V., VARKI A., HENRIQUES-NORMARK B.  
*Streptococcus pneumoniae* senses a human-like sialic acid profile via the response regulator CiaR.  
*Cell Host Microbe*. 2016 Sep 14;20(3):307-17.
- II. **HENTRICH K.**, SENDER V., PATHAK A., HENRIQUES-NORMARK B.  
Human sialic profiles mediate increased pneumococcal adhesion and immune evasion.  
*Manuscript*
- III. SENDER V., **HENTRICH K.**, PATHAK A., NORMARK S., HENRIQUES-NORMARK B.  
Mechanism for enhanced bacterial burden in the lower respiratory tract of mice during influenza/pneumococcal coinfection.  
*Manuscript*
- IV. SPELMINK L., SENDER V., **HENTRICH K.**, KURI T., PLANT L., HENRIQUES-NORMARK B.  
Toll-like receptor 3/TRIF-Dependent IL-12p70 secretion mediated by *Streptococcus pneumoniae* RNA and its priming by influenza A virus coinfection in human dendritic cells.  
*MBio*. 2016 Mar 8;7(2):e00168-16.



## **Publications by the author, which are not included in the thesis**

SCHULTE T., LÖFLING J., MIKAELSSON C., KIKHNEY A., **HENTRICH K.**, DIAMANTE A., EBEL C., NORMARK S., SVERGUN D., HENRIQUES-NORMARK B., ACHOUR A.

The basic keratin 10-binding domain of the virulence-associated pneumococcal serine-rich protein PsrP adopts a novel MSCRAMM fold.

*Open Biol.* 2014 Jan 15;4:130090.

ORRSKOG S., ROUNIOJA S., SPADAFINA T., GALLOTTA M., NORMAN M., **HENTRICH K.**, FÄLKER S., YGBERG-ERIKSSON S., HASENBERG M., JOHANSSON B., UOTILA L.M., GAHMBERG C.G., BAROCCHI M., GUNZER M., NORMARK S., HENRIQUES-NORMARK B.

Pilus adhesin RrgA interacts with complement receptor 3, thereby affecting macrophage function and systemic pneumococcal disease.

*MBio.* 2012 Dec 26;4(1):e00535-12.

MONTEIRO C., PAPENFORT K., **HENTRICH K.**, AHMAD I., LE GUYON S., REIMANN R., GRANTCHAROVA N., RÖMLING U.

Hfq and Hfq-dependent small RNAs are major contributors to multicellular development in *Salmonella enterica* serovar Typhimurium.

*RNA Biol.* 2012 Apr;9(4):489-502.



# CONTENTS

1	Introduction.....	1
1.1	<i>Streptococcus pneumoniae</i> .....	1
1.1.1	Pneumococcal colonization and disease .....	2
1.1.2	Treatment and prevention of pneumococcal infections.....	5
1.2	The immune system .....	8
1.2.1	The innate immune response.....	8
1.2.2	The adaptive immune response.....	13
1.3	Pneumococcal-Host interactions.....	14
1.3.1	Host glycans and their biological functions.....	14
1.3.2	Pneumococcal virulence factors.....	20
2	Aims.....	27
2.1	Specific aims .....	27
3	Methodological considerations .....	29
3.1	Bacterial and viral strains and their growth conditions .....	29
3.2	Characterization of pneumococcal phenotypes .....	30
3.2.1	Sialidase activity.....	30
3.2.2	Production of hydrogen peroxide and acetyl phosphate.....	30
3.2.3	Survival in response to hydrogen peroxide.....	30
3.3	In vitro and ex vivo models.....	31
3.3.1	Cells .....	31
3.3.2	Sialic acid feeding of cells.....	31
3.3.3	Measurement of cytokine response and cell toxicity.....	32
3.4	In vivo models.....	32
3.5	Ethical considerations .....	32
4	Results and Discussion .....	33
4.1	Paper I.....	33
4.2	Paper II .....	36
4.3	Paper III .....	39
4.4	Paper IV.....	42
5	Concluding remarks.....	44
6	Acknowledgements .....	45
7	References.....	47

## LIST OF ABBREVIATIONS

AcPh	Acetyl phosphate
AIM2	Absent in melanoma 2
AMs	Alveolar macrophages
AMPs	Antimicrobial peptides
AOM	Acute otitis media
ASC	Apoptosis-associated speck-like protein containing CARD
Asn	Asparagine
BAL	Bronchoalveolar lavage
CAP	Community-acquired pneumonia
CARD	Caspase recruitment domain
CD	Cluster of differentiation
CD33r Siglec	CD33-related Siglec
CFU	Colony-forming units
CMAH	CMP-Neu5Ac hydroxylase
CO <sub>2</sub>	Carbon dioxide
CR	Complement receptor
CRAMP	Cathelicidin-related antimicrobial peptide
DCs	Dendritic cells
DNA	Deoxyribonucleic acid
dsRNA	Double-stranded RNA
Fuc	Fucose
Gal	Galactose
GalNAc	<i>N</i> -acetylgalactosamine
GBS	Group B streptococci
Glc	Glucose
GlcNAc	<i>N</i> -acetylglucosamine
hBD	Human $\beta$ -defensin
HIV	Human immunodeficiency virus
HK	Histidine kinase
HNP	Human neutrophil peptides
HtrA	High temperature requirement A
H <sub>2</sub> O <sub>2</sub>	Hydrogen peroxide
IAV	Influenza A virus
IFN	Interferon
Ig	Immunoglobulin
IL	Interleukin
IPD	Invasive pneumococcal disease
IRF	Interferon regulatory factor
LPS	Lipopolysaccharide
LRT	Lower respiratory tract
LTA	Lipoteichoic acid
MAAII	<i>Maackia amurensis</i> lectin II
Man	Mannose
ManNAc	<i>N</i> -acetylmannosamine
MARCO	Macrophage receptor with collagenous structure
MHC	Major histocompatibility complex

MIP	Macrophage inflammatory protein
MOI	Multiplicity of infection
MyD88	Myeloid differentiation factor 88
NETs	Neutrophil extracellular traps
Neu5Ac	<i>N</i> -acetylneuraminic acid
Neu5Gc	<i>N</i> -glycolylneuraminic acid
NF- $\kappa$ b	Nuclear factor $\kappa$ b
NLR	NOD-like receptor
NOD	Nucleotide-binding oligomerization domain
PAMPs	Pathogen associated microbial patterns
PBP	Penicillin-binding protein
PCR	Polymerase chain reaction
PCV	Pneumococcal conjugate vaccine
PGN	Peptidoglycan
Ply	Pneumolysin
PPSV	Pneumococcal polysaccharide vaccine
PRR	Pattern recognition receptor
RNA	Ribonucleic acid
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
RR	Response regulator
Ser	Serine
Sia	Sialic acid
Siglec	Sialic acid binding Ig-like lectins
SNA	<i>Sambucus nigra</i> lectin
SP-A, SP-D	Surfactant protein-A, Surfactant protein-D
SpxB	Streptococcal pyruvate oxidase B
STING	Stimulator of interferon genes
TCS	Two-component system
T <sub>H</sub> -cells	Helper T-cells
Thr	Threonine
TIR	Toll-interleukin 1 (IL-1) receptor
TLR	Toll-like receptor
TNF $\alpha$	Tumor necrosis factor $\alpha$
TRIF	TIR-domain-containing adapter protein-inducing interferon- $\beta$
WT	Wild-type




# 1 INTRODUCTION

## 1.1 *Streptococcus pneumoniae*

In 1881, George Miller Sternberg and Louis Pasteur isolated diplococci after injecting human saliva into rabbits [1, 2]. In 1886, Fraenkel used the name *Pneumococcus* for the first time, referring to the pneumonia these bacteria cause. In the same decade, Christian Gram developed a staining to visualize bacteria in patient samples. He discovered lancet-shaped gram-positive cocci in the lung specimens of patients who died of pneumonia and named them “the cocci of croupous pneumonia”. In 1920, the isolate was given the name *Diplococcus pneumoniae*, due to its microscopic morphology, which was seen in the gram-stain. It took until 1974 that these bacteria were referred to as *Streptococcus pneumoniae* [3]. However, the term “*Pneumococcus*” is still frequently used.

Pneumococcal colonies appear to be small and greyish on blood agar plates after an overnight incubation at 37°C and 5% carbon dioxide (CO<sub>2</sub>). Moreover, they are  $\alpha$ -hemolytic, as they oxidize hemoglobin. To date, we know of at least 97 different pneumococcal serotypes, which differ in the composition of their polysaccharide capsule [4]. The presence of the capsule leads to the mucoid appearance of pneumococcal colonies on solid agar. Another characteristic of most *S. pneumoniae* isolates is the sensitivity to optochin, which allows differentiating them from *S. viridans* [5, 6] (Figure 1). However, this test is not sufficient to identify pneumococci, since there have been recent reports of optochin-resistant isolates [7].




**Figure 1** Optochin sensitivity of the  $\alpha$ -hemolytic *S. pneumoniae* TIGR4. TIGR4 was streaked out on a blood agar plate and cultured at 37°C in the presence of 5% CO<sub>2</sub>. The sensitivity to optochin was tested with a disk diffusion method.

*S. pneumoniae* was the first bacterium shown to be naturally competent. In 1928, Griffith showed that viable non-virulent pneumococci turned into virulent ones when they were injected into mice together with heat-killed pathogenic isolates. He hypothesized that the dead pneumococcal strain provides a certain protein that enables the attenuated bacteria to be virulent [8]. It took 16 more years to show that the exchanged substance, responsible for the increased virulence, was not protein, but deoxyribonucleic acid (DNA) [9].

### 1.1.1 Pneumococcal colonization and disease

Pneumococci are human-adapted and often found asymptotically colonizing the upper respiratory tract. Nonetheless, *S. pneumoniae* can cause diseases ranging from benign respiratory tract infections, e.g. sinusitis or otitis media, to severe diseases like pneumonia, septicaemia and even meningitis [10], making pneumococcal infections a major cause of global childhood mortality, leading to about 11% of all deaths among children below the age of five [11] (Figure 2).



**Figure 2 Pneumococcal mortalities of children younger than 5 years.** The graph represents the number of deaths per 100 000 HIV-negative children caused by pneumococcal infections. The picture was adopted from [11].



## ***Colonization***

Pneumococcal colonization is believed to occur via the spread of pneumococci-containing aerosols. Before their eventual clearance by the immune system, pneumococci can colonize the upper respiratory tract for several weeks or months [12]. Especially pre-school age children and those attending day care centers are found to be the main reservoir for *S. pneumoniae* [13, 14], with carriage rates up to 60% [10, 15]. Less than 10% of adults are carriers, although a higher colonization rate was found in parents of small children [16]. Once these bacteria establish colonization, they can replicate and form biofilms and/or translocate to other sites and cause disease.

## ***Otitis media and sinusitis***

*S. pneumoniae* is one of the causative agents of acute otitis media (AOM), an inflammation of the middle ear, affecting up to 85% of all children at the age of 3 years [17]. In Sweden and the US, AOM represents the most common infectious disease to prescribe antibiotics to children [18]. However, recurrent or persistent AOM can lead to complications and severe sequelae, e.g. conductive hearing loss, labyrinthitis, perforation of the tympanic membrane or even meningitis [19].

Although viruses cause most of the sinusitis infections, *S. pneumoniae* is among the most common bacterial agents that can lead to infections of the paranasal sinus cavity [20]. While most viral sinusitis episodes resolve completely after 10 days, bacterial rhinosinusitis is usually more persistent [21].

## ***Pneumonia***

Pneumococcal infections of normally sterile sites of the body, like the lower respiratory tract, the blood or meninges, are considered invasive pneumococcal diseases, also called IPD [22]. An inflammation of the lungs is called pneumonia. *S. pneumoniae* is the most common cause of community-acquired pneumonia (CAP), followed by *Haemophilus influenzae*, respiratory viruses or *Mycoplasma pneumoniae* [23]. CAP represents a huge financial and clinical burden throughout the whole world [24] and accounts for 19% of all mortality cases in children below the age of 5 years. The distribution of pneumonia-caused deaths differs greatly between geographical regions and their economic situation, with 50% of all pneumonia-related mortalities occurring in the African and only 2% - 3% in the European and American region [25].

### ***Bacteraemia and sepsis***

20-30% of the children with pneumonia develop bacteraemia, the occurrence of bacteria in the blood stream [26]. The consequence can be sepsis, a life-threatening systemic infection. CAP and health-care associated pneumonia in children and adults are the pre-conditions, which lead to 50% of all sepsis cases. Especially, very young children below 1 year of age and elderly people of more than 65 years of age are prone to develop sepsis [27].

### ***Meningitis***

Since the introduction of the vaccine against *H. influenzae* type b, *S. pneumoniae* and *Neisseria meningitidis* present the major causes of bacterial meningitis in children nowadays [26, 28]. A study conducted in the US, observed that mortality rates in children were less than 10%, but almost half of the surviving patients suffered from severe sequelae, like hearing loss, brain infarcts, brain abscesses or hydrocephalus [29]. European studies describe a fatality rate of 30% in adults after pneumococcal meningitis and cognitive deficits in 26% of the survivors [30, 31]. As it is the case for pneumonia, the mortality rates after pneumococcal meningitis are significantly higher in developing than in developed countries [11].

### ***Risk factors***

As discussed before, the age determines the risk of IPD, affecting especially very young children with their naïve immune system, elderly people of an age of more than 65 years due to their weakening immune response, and individuals with co-morbidities, e.g. diabetes or different immunodeficiencies [32]. Environmental and social factors, such as the number of siblings, financial income, and smoking affect pneumococcal colonization. Furthermore, the ethnic background was also shown to determine the risk of a pneumococcal infection, with African Americans and Native Americans having a high probability of developing IPD [33].

Coinfections with viral pathogens, especially with influenza A virus (IAV, see below) and human immunodeficiency virus (HIV), have been shown to significantly promote pneumococcal infections [32, 34].

### ***Pneumococcal coinfection with Influenza A virus***

During influenza pandemics, most fatalities are not caused by the virus alone, but by a secondary infection with a bacterial pathogen [35]. In 1918, the “Spanish flu” was caused by subtype H1N1. This pandemic killed more than 50 million people, which primarily died due to superinfections with *S. pneumoniae* [36]. In 1957, IAV subtype H2N2 caused the “Asian flu”, leading to more than 1 million deaths partially due to pneumonia caused by *Staphylococcus aureus*. The severe and about 500 000 fatal cases of the “Hong Kong flu” in

1968 were mainly caused by co-infections of influenza subtype H3N2 and pneumococci [37-39]. The next influenza pandemic occurred 41 years later, and is known as the so-called “swine flu”. In 2009, infections with H1N1 led to about 200 000 deaths with mainly young adults to be affected. Like former pandemics, severe cases and fatalities were mainly caused by *S. aureus* and *S. pneumoniae* [40]. Although it was first believed that the introduction of antibiotics against *S. pneumoniae* was responsible for the shift from secondary infections with pneumococci to co-infection with other pathogens, strain-related changes in the virus and/or the bacteria are more probable to be responsible for variations in the severity of influenza pandemics. Additionally, the introduction of the pneumococcal vaccine is also believed to account for lower numbers of coinfections with *S. pneumoniae* during the pandemic in 2009 [35].

### 1.1.2 Treatment and prevention of pneumococcal infections

#### *Treatment*

The treatment of choice of pneumococcal infections is a therapy with antibiotics. Since the introduction in the 1940s, penicillin was used against infections with *S. pneumoniae*.  $\beta$ -lactam-antibiotics, like penicillin, inhibit the bacterial cell-wall synthesis by interfering with penicillin-binding proteins (PBPs). Mutations in several different genes, e.g. pneumococcal PBPs or the cell-wall mucopeptide branching enzyme MurM, are shown to promote resistance development to penicillins [41]. Already in the 1960s, reports about  $\beta$ -lactam-resistant pneumococci were published [42], and in 2013, Sweden reported that 6.8% of all IPD isolates were non-susceptible to penicillins [43].

Following the isolation of increasing numbers of penicillin-resistant pneumococci, the use of macrolides and fluoroquinolones has become more common to treat respiratory tract infections. Macrolides, like erythromycin or the semi-synthetic azithromycin, inhibit bacterial protein synthesis by binding to the 50S ribosomal subunit and inhibit protein elongation [44]. Fluoroquinolones, e.g. moxifloxacin, are synthetically produced antibiotics that bind to type II topoisomerase enzymes, like DNA gyrase, thereby preventing DNA replication and cell division. Macrolide-resistant pneumococci have been isolated in several countries with resistance rates up to 80% [45]. In 2013, 6.5% of invasive pneumococci, isolated in Sweden, were non-susceptible to macrolides [43]. A study from the same year, conducted in the US, reported that about one third of all cases of IPD were caused by pneumococci, which were resistant to several antibiotics [46].

## Prevention

In order to prevent the spread of antibiotic-resistant pneumococci and the development of IPD, a pneumococcal vaccine was introduced. To date, there are 4 different pneumococcal vaccines available: A pneumococcal polysaccharide vaccine (PPSV23) and pneumococcal conjugate vaccines (PCV7, PCV10 and PCV13), which are shown to protect against the most common pneumococcal serotypes linked to IPD (Table 1).

**Table 1 Pneumococcal vaccines currently available [47]**

Name	Supplier	Licensed	Serotypes covered
PPSV23 Pneumovax®23	Merck	1983	1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, 33F
PCV7 Prevnar®/ Prevenar®	Pfizer	2000	4, 6B, 9V, 14, 18C, 19F, 23F
PCV10 Synflorix™	GSK	2009	1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F
PCV13 Prevenar 13®	Pfizer	2010	1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F

In 1983, a pneumococcal vaccine covering 23 different serotypes was licensed. This vaccine is polysaccharide-based and provokes an antibody response against the polysaccharide capsule of pneumococci, which promotes opsonophagocytosis. However, polysaccharide-based vaccines elicit no or only weak immune responses in very young children and adults and fail to induce immune memory due to their T-cell independent response (see chapter 1.2.2) [48].

In 2000 and 2001, the first PCV was licensed in the US and Europe, respectively [49]. PCVs consist of 7, 10 or 13 different capsular polysaccharides, which are coupled to protein carriers, a non-toxic variation of the diphtheria toxin, CRM197 (PCV7 and PCV13) or protein D from *H. influenzae* (PCV10). These vaccines cause T-cell dependent immune responses and B-cell memory [50]. The introduction of PCVs into national vaccination programmes in Europe decreased pneumococcal infections drastically in vaccinated individuals, but also in non-vaccinated people because of herd-immunity. In Germany, the introduction of PCV7 was recommended from 2006 onwards and led to a vaccine-coverage of 84% in 2007. In 2008, Germany reported a 50% reduction in IPD cases in children below the age of 2 years [51]. Sweden incorporated PCVs into the national immunization programme in 2009. Of all children born in 2013 in Sweden, 96.6% were vaccinated against pneumococcal infections by the age of 2 years [52], which led to a considerably reduction of IPD cases in vaccinated children [53].

However, although PCVs drastically reduce the nasopharyngeal carriage rates and IPD cases caused by pneumococcal serotypes included in the vaccines, other serotypes, not included in the vaccine, are found to replace the serotypes in colonization and pneumococcal disease [54, 55]. In Sweden, 36% of the children between 1 and 5 years were colonized with pneumococci before the introduction of pneumococcal vaccines. After the pneumococcal vaccines were included into the childhood immunisation programme, the carriage rate was not severely different with 30%, which is attributed to serotype replacement [53].

In order to produce a vaccine that targets all pneumococcal serotypes simultaneously, several studies are concentrating on virulence factors present in all isolates. For example, the pneumococcal surface protein A (PspA), the pneumococcal surface antigen A (PsaA), as well as pneumolysin have been shown to mediate immunity against pneumococcal infections (See chapter 1.3.2) [56-58]. Moreover, immunization experiments with the pneumococcal surface protein C (PspC) were also shown to protect against colonization [59]. Increased protection rates have been observed, if these antigens were combined in immunization studies [57].

## 1.2 The immune system

The function of the immune system is to distinguish “self” from “non-self” and evading pathogens from non-pathogens, in order to prevent an infection, remove tumour cells and maintain an immune homeostasis. The immune system consists of lymphatic organs, physical barriers, cells and soluble mediators and can be divided into the innate and the adaptive immune response.

The innate immune system comprises epithelial barriers, e.g. skin and mucosa in the respiratory or gastrointestinal tract, cells, like macrophages, neutrophils or dendritic cells (DCs), and soluble components, like the complement system and antimicrobial peptides (AMPs). It is the first line of defence during an infection, with a rapid and non-specific response that is not adapted to the type of pathogen. Moreover, it regulates the adaptive immune response, e.g. by producing cytokines. In contrast to the innate immune response, the adaptive response is highly specific, adjusted to the pathogen and able to develop an immunological memory. Its function is mediated by B- and T-lymphocytes and soluble factors, like antibodies.

### 1.2.1 The innate immune response


#### *Detection of invading pathogens and production of inflammatory mediators*

The innate immune system is activated by pathogen-associated molecular patterns (PAMPs), structures, which are part of microbial pathogens and are required for their virulence. Usually, they are common to many different pathogens. Examples of PAMPs are bacterial DNA and lipopolysaccharide (LPS) [60]. PAMPs bind to and activate pattern-recognition receptors (PRR) on or inside host cells, e.g. Toll-like receptors (TLRs), nucleotide-binding oligomerization domain (NOD)-like receptors (NLRs) or cytosolic DNA receptors.

To date, we know of 10 human and 13 murine TLRs. While TLR1, TLR2, TLR4, TLR5, TLR6 and TLR11 are positioned on the plasma membrane, TLR3, TLR7, TLR8 and TLR9 are located in endosomes. TLRs consist of an ectodomain, required for the recognition of PAMPs, transmembrane domains, and Toll-interleukin 1 (IL-1) receptor (TIR) domains. There are different adapter molecules, known to interact with the TIR domain of TLRs, and which are required to evoke a response by the host cell. All TLRs, except for TLR3, known to recognize double-stranded ribonucleic acid (dsRNA), use the myeloid differentiation factor 88 (MyD88) as an adapter molecule to induce the transcription of inflammatory cytokines by activating the transcription factor nuclear factor-kappa B (NF- $\kappa$ B) and mitogen-activated protein kinases (MAPKs). TLR3, and also TLR4, use a TIR-domain-containing adapter

protein-inducing interferon- $\beta$  (TRIF) as an adapter molecule to stimulate the transcription factors interferon regulatory factor 3 (IRF-3) and NF- $\kappa$ B, leading to the induction of type I interferon (IFN) and inflammatory cytokines, like interleukin-6 (IL-6) or tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) [61, 62].

NLRs are located in the cytoplasm of cells. To date, we know of at least 22 different types in humans, and 33 in mice. NLRs consist of C-terminal leucine-rich repeats that recognize PAMPs, a nucleotide-binding oligomerization domain (NOD) and a variable N-terminal protein-protein interaction domain, e.g. caspase recruitment domain (CARD), that is crucial for inducing downstream signals, like NF- $\kappa$ B activation, leading to the transcription of pro-inflammatory cytokines [63]. Inflammasomes, consisting of nod-like receptor protein 3 (NLRP3) and absent in melanoma 2 (AIM2), are intracellular protein complexes that are activated upon bacterial infections. They recruit apoptosis-associated speck-like protein containing CARD (ASC), which binds to caspase-1 and evokes cleavage of the pro-forms of IL-1 $\beta$  and IL-18, thus leading to their maturation [62, 64].



**Figure 3 Schematic representations of selected PRRs and their known pneumococcal PAMPs.** Pneumococcal LTA is sensed by TLR2, while TLR4 has been proposed to recognize Ply. The endosomal TLR9 senses DNA. TLRs that use MyD88 as an adapter (TLR2, TLR4, TLR9) induce the transcription of inflammatory cytokines, TLR3 and TLR4 signal via TRIF and thus activate additionally the expression of type I IFN. Ply is able to activate the NLRP3/AIM2 inflammasome and hence leads to the production of IL-1 $\beta$ . PGN activates NOD2 and the transcription of inflammatory cytokines, while DNA that signals via STING additionally induces type I IFN.

Various pneumococcal structures can be sensed by PRRs. Lipoteichoic acid (LTA), a component of the pneumococcal cell wall, is sensed by TLR2 [65]. Unmethylated CpG motifs in pneumococcal DNA can be recognized by TLR9 [66], and the pneumococcal toxin pneumolysin (Ply) has been suggested to be sensed by TLR4 [67]. Peptidoglycan (PGN), another pneumococcal cell wall component, can be sensed by NOD2, dependently on the activity of the pneumococcal toxin pneumolysin [68, 69]. Moreover, pneumolysin is capable of activating the inflammasome and subsequent production of IL-1 $\beta$  and IL-18 [70, 71] and is also required for the pneumococcal activation of the cytosolic DNA receptor stimulator of interferon genes (STING) [72] (Figure 3).

### ***Macrophages***

Bone marrow-derived monocytes, which circulate in the blood, are recruited and differentiate into macrophages and DCs during an inflammation [73]. Macrophages play a central role in the immune response and are present in many tissues and organs, especially those, which are exposed to the environment [74]. In the lungs, residential alveolar macrophages (AMs) represent 95% of all cells, and are therefore the major cell type [75].

The key function of macrophages is the phagocytosis and killing of particles and pathogens, both opsonized and non-opsonized [75, 76]. Invading microbes can be recognized and taken up into the cell due to binding to different receptors on the cell surface, like C-type lectins (e.g. mannose receptor), scavenger receptors (e.g. macrophage receptor with collagenous structure (MARCO)), complement receptors (CR) or immunoglobulin receptors (Fc $\gamma$  receptors) [77, 78].

Following phagocytosis, pathogens are eradicated by reactive oxygen species (ROS) or reactive nitrogen species (RNS), e.g. superoxide, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) or nitric oxide [79]. Furthermore, host cells secrete pro-inflammatory cytokines and chemokines, like IL-8, which recruit neutrophils to the site of infection, and the production of monocyte chemoattractant protein-1 (MCP-1) or RANTES evokes an influx of activated monocytes and lymphocytes [75].

Macrophages play a significant role in fighting pneumococcal pneumonia and the importance of several receptors in pneumococcal uptake has been described in the literature. Scavenger receptors, such as MARCO [80], C-type lectins, such as SIGN-R1 [81], or the complement receptor CR3 [82] have been shown to mediate pneumococcal phagocytosis. Pneumococci, on the other hand, evade phagocytosis mechanisms by the host with the help of their thick polysaccharide capsule [83]. Although the production of ROS has been implicated in macrophage-mediated killing of different bacteria, pneumococcal clearance by macrophages is facilitated by RNS, and not attributed to ROS production [84].



## ***Neutrophils***

Neutrophils are produced in the bone marrow and are afterwards released into the vascular system, where they represent 60% of all leukocytes in humans [85]. During inflammation in the lungs, macrophages and epithelial cells produce cytokines and chemokines, which recruit neutrophils into the alveolar space [86, 87]. Like macrophages, neutrophils phagocytose bacteria and eradicate them with the help of ROS, but also non-oxidative mechanisms, like AMPs, proteases and the formation of neutrophil extracellular traps (NETs) [88, 89].

As mentioned above, *S. pneumoniae* was shown to avoid phagocytosis and to be resistant to ROS. Instead, neutrophils fight pneumococcal infections with serine proteases [90, 91]. In order to escape killing by neutrophils, pneumococci have developed different immune evasion strategies. The pneumococcal capsule was shown to promote escape from NETs [92], and pneumococcal endonuclease A degrades DNA, the main building block of NETs [93].

## ***Dendritic cells***

DCs account for only 1% of all immune cells [94]. They originate in the bone marrow, and represent the connection between the innate and the adaptive immune response. These cells express a number of receptors, like scavenger, complement receptors or Fc receptors, which promote phagocytosis and are thus required for antigen presentation, the most important function of DCs [95].

DCs are located in the mucosa, where they constantly collect antigens. After maturation and translocation to the lymph nodes, they present these antigens to T-cells via major histocompatibility complex class II (MHC class II) molecules, leading to the activation of T-cells and the initiation of an adaptive immune response [96].

## ***Antimicrobial peptides and collectins***

AMPs are effective against Gram-positive and Gram-negative bacterial, viral, as well as fungal pathogens and show a great variation in size (6 - 59 amino acids), sequence and secondary structure. They bind to microbes and kill them either by disturbing membrane integrity and causing lysis, or they enter the cell and interfere with essential mechanisms, e.g. nucleic acid synthesis [97].

In the respiratory tract of humans, primarily epithelial cells and neutrophils produce AMPs, which help to kill invading pathogens [98]. Neutrophil  $\alpha$ -defensins / human neutrophil peptides (HNPs), human  $\beta$ -defensins (hBDs), and cathelicidins, like hCAP18/LL-37 are the major AMPs present in the lungs and bronchoalveolar lavage (BAL) (see Table 2) [99].

**Table 2 AMP-producing cells in the respiratory tract of humans [99, 100]**

Cell type	AMPs	Function
Neutrophils	$\alpha$ -defensins (HNP 1-4)	Microbial killing
Epithelial cells	hBDs	Microbial killing Chemo-attraction of immune cells Activation of DCs
Neutrophils Epithelial cells	hCAP18/LL-37	Microbial killing

Some hBDs, like hBD-1, are constitutively expressed, while other hBDs are induced e.g. by TLR-signalling or cytokine expression [100]. LL-37 is shown to be resistant against proteolytic activity due to its ability to form aggregates [101]. Its expression is regulated by inflammatory signalling, Vitamin D and endoplasmic reticulum stress. The murine homologue of LL-37 cathelicidin-related antimicrobial peptide (CRAMP) [102] and several defensins have been identified in mice [103], which are suggested to play an important role in murine host defence mechanisms in the lung [104].

Pneumococci are sensitive to treatment with LL-37 from mast cells [105] and human  $\alpha$ -defensins produced by neutrophils [106]. Furthermore, primary human lung epithelial cells have been shown to secrete hBD-2 and hBD-3 in response to pneumococcal infections, which promoted bacterial clearance [107].

In the lungs, surfactants are secreted by type II alveolar epithelial and Clara cells [108] and consist of 90% lipids and 10% proteins. They play an important role in reducing surface tension, and take part in the innate immune response. There are several surfactant proteins (SP). This thesis includes SP-A and SP-D, which are hydrophilic and belong to the group of collagen-containing C-type lectins (carbohydrate-binding proteins), so-called collectins. SP-A is believed to stabilize surfactants, while SP-D is involved in keeping the homeostasis. Moreover, both proteins opsonize pathogens, promoting their clearance [109, 110].

Recombinant SP-D was found to bind to and agglutinate pneumococci, but did not promote neutrophil-mediated killing [111]. Using an *in vivo* model of intranasal challenge, it was demonstrated that SP-D endorses pneumococcal clearance and prevents bacterial translocation from the upper to the lower respiratory tract [112]. SP-A, on the other hand, was shown to promote pneumococcal uptake by alveolar macrophages via the scavenger receptor [113].

## 1.2.2 The adaptive immune response

### *T-lymphocytes*

Naïve T-lymphocytes (or T-cells) originate in the thymus and are found in the blood and lymphatic organs. T-cell receptors on their surface recognize antigens which are presented via MHC molecules, e.g. by DCs. This leads to the activation of the T-cells, evoking either a cytotoxic function or the production of cytokines. Cytotoxic T-cells (or CD8<sup>+</sup> T cells) are activated by MHC class I molecules and have the ability to directly kill infected cells, e.g. by lysis. CD4<sup>+</sup> helper T-cells, on the other hand, are activated by MHC class II molecules and produce cytokines, which can be lethal to infected cells, and lead to immunoglobulin (Ig) production by B-cells or stimulate other T-cell functions [114].

After an infection with *S. pneumoniae*, DCs produce large amounts of IL-12, a pro-inflammatory cytokine that promotes the differentiation of helper T-cells (T<sub>H</sub>-cells). Several studies have elucidated the importance of IFN- $\gamma$  secretion by T<sub>H</sub>-1 cells in *in vivo* models after pneumococcal challenge [115-117].

### *B-lymphocytes*

B-lymphocytes (or B-cells) mature in the bone marrow and play a major role in the adaptive immune response. B-cells bind and take up antigens, in order to present them on their surface by MHC class II molecules. These antigens can now be recognized by specific T<sub>H</sub>-cells, which in turn stimulate the differentiation of B-cells into plasma cells producing high-affinity class-switched antibodies (IgG, IgE or IgA) or memory B-cells [118, 119].

Carbohydrates alone, like the pneumococcal vaccine PPSV23, activate B-cells in a T-cell independent manner, since they cannot be presented by MHC class II. Therefore, B-cells are only capable of producing short-lived low-affinity antibodies, like IgM. Subsequently, the immune response does not last long. Furthermore, children under the age of 2 years are not able to evoke an immune response to carbohydrates only, since their B-cells are not fully matured. They hardly express type 2 complement receptors and lack certain cytokines, needed to activate B-cells. In order to induce an immune response in very young children, the pneumococcal vaccines PCV7, PCV10 and PCV13 were designed differently. The pneumococcal polysaccharides have been linked to a carrier protein, which can be presented by MHC class II molecules and this leads to a T-cell dependent immune response [118].

### 1.3 Pneumococcal-Host interactions

Carbohydrates are essential structures to enable pneumococcal interactions with the host. They facilitate pneumococcal colonization and invasive disease but also evoke a host immune response. The following chapter will discuss host glycosylation, with an emphasis on sialic acids (Sias), the carbohydrates that are found on the tip of most glycan strands. In order to hide and escape from the host's immune system, pneumococci comprise several virulence factors, of which the pneumococcal capsule, the sialidase NanA, the main Sia transporter SatABC, the two-component system CiaRH, the streptococcal pyruvate oxidase SpxB, the serine protease HtrA, and pneumolysin will be described. Moreover, the pathogenesis of influenza/pneumococcal coinfections will be discussed.

#### 1.3.1 Host glycans and their biological functions


##### *Host glycosylation*

In 1967, Rambourg and colleagues observed that the cell surface is highly covered with sugar molecules, the glycocalyx [120]. Glycans exist in many variations and are built of mono- or oligosaccharides, which are covalently linked to a non-carbohydrate moiety, a protein or lipid, or exist freely [121]. The most common monosaccharides present in mammalian glycoconjugates are listed in Table 3.

**Table 3 The most common monosaccharides in mammalian glycans [121]**

Name	Structure	Examples
Sialic acids	Nine-carbon backbone acidic sugar	<i>N</i> -acetylneuraminic acid, <i>N</i> -glycolylneuraminic acid
Hexoses	Six-carbon neutral sugars	Glucose, galactose, mannose
Hexosamines	Amino group at position 2 of an hexose which is free or <i>N</i> -acetylated	<i>N</i> -acetylglucosamine, <i>N</i> -acetyl-galactosamine
Deoxyhexoses	Six-carbon neutral sugar without hydroxylgroup at position 6	Fucose
Pentoses	Five-carbon sugar	Xylose
Uronic acids	Hexose with negatively charged carboxylate at position 6	Glucuronic acid, Iduronic acid

Glycosylation is the most common post-translational modification of proteins. Depending on their linkage to a protein, the aglycone, glycans are mainly divided into N- and O-linked. N-linked glycans are covalently attached to a polypeptide via an asparagine (Asn) residue. O-linked glycans are linked to the polypeptide via *N*-acetylgalactosamine (GalNAc) on a serine (Ser) or threonine (Thr) residue (Figure 4). In eukaryotes, these glycosylations take place in the cytoplasm, the endoplasmatic reticulum and golgi [121, 122].



**Figure 4 Schematic representations of common mammalian glycans.** Glycoproteins and glycolipids are usually decorated with chains of GalNAc, galactose (Gal), *N*-acetylglucosamine (GlcNAc), mannose (Man), fucose (Fuc) or glucose (Glc) molecules, with a terminal Sia molecule. Modified after Varki (2007) [123].

Other life forms such as plants and bacteria differ greatly from mammals in their oligosaccharide composition. To date, several glycans are exclusively found in bacteria, e.g. 3-deoxy-D-manno-oct-2-ulonic acid (KDO), presenting part of the core of the bacterial endotoxin LPS, and *N*-acetylmuramic acid (MurNAc), found in peptidoglycan, the main building block of the bacterial cell wall [124].

### ***Sialic acids***

The biochemists Blix and Klenk discovered Sias in glycolipids of the brain and in salivary mucins [125, 126]. The terminal monosaccharide of glycoconjugates, like glycoproteins and glycolipids, is usually a nine-carbon backbone sugar, called neuraminic acid or Sia, with

modifications mainly occurring at position 4, 5, 7, 8 and 9 [121, 127]. Sias are found on all cells in animals of the deuterostome lineage (from sea urchins to mammals) [123, 128], and in some bacteria, fungi and protozoa [129]. Although Sias are usually not present in insects, they have been detected in early developmental stages of *Drosophila melanogaster* and the cicada *Philaenus spumarius* [130, 131].

To date, there are about 50 different types of Sia known in nature. Neu5Ac (*N*-acetylneuraminic acid) and Neu5Gc (*N*-glycolylneuraminic acid) are the most common ones [128]. Most of them are attached by their 2-carbon to the 3- or 6-carbon of Gal or GalNAc underneath, commonly referred to as  $\alpha$ 2-3- or  $\alpha$ 2-6-linkage [121].

### ***CMP-NeuAc hydroxylase (CMAH)***


In most animals, the enzyme CMP-Neu5Ac hydroxylase (CMAH) transforms CMP-Neu5Ac into CMP-Neu5Gc by adding a single oxygen atom [132]. About 3 million years ago, a 92-bp deletion in the exon 6 encoding *CMAH* occurred in humans. Consequently, humans are only able to synthesize Neu5Ac, while many other mammals, including our closest relatives the chimpanzees, present mainly Neu5Gc as terminal sugar on their glycoconjugates (Figure 5) [133, 134]. The occurrence of the *CMAH* mutation was predicted to have happened when stone tools were used to butcher hunted animals, which might have caused minor injuries and infections [135]. It was suggested that evolutionary pressure caused by Neu5Gc-recognizing pathogens and higher fertility of Neu5Gc-negative females led to the positive selection of CMAH-negative individuals [136, 137].

Noteworthy, humans are not the only species with a non-functional CMAH. Recently, it was described that New World Monkeys obtained a mutation in their *CMAH* gene already 30 million years ago [138]. Likewise, ferrets have a mutation in *CMAH* in their genome, which was shown to increase their susceptibility to human IAV [139].

Different human-adapted pathogens have evolved their virulence mechanisms in accordance to the human Sia profile. The merozoite of *Plasmodium falciparum* recognizes Neu5Ac, but not Neu5Gc [140], therefore infecting New World monkeys but no other primates [141]. Similarly, the typhoid toxin of *Salmonella Typhi* was shown to strongly bind to Neu5Ac, but only weakly to Neu5Gc [142].

The loss of Neu5Gc is shown to affect B-cell reactivity in *Cmah*<sup>-/-</sup> mice, since its absence leads to enhanced levels of B-cell proliferation and antibody production [143]. Similarly, T-cells of *Cmah*<sup>-/-</sup> mice exhibit a greater activation upon stimulation compared to wt cells, which can be repressed by Neu5Gc-treatment [144]. Moreover, in comparison to chimpanzees, humans display a more active immune system. Both B- and T-lymphocytes of humans are reported to react stronger to stimuli than those of chimpanzees [145].

Although humans are not able to synthesize Neu5Gc, this Sia can be incorporated into human tissues through their diet. Neu5Gc is antigenic in humans, evoking an antibody response [146], which was suggested by Hedlund *et al.* to promote inflammation and the development of cancer [147] and later experimentally confirmed by Samraj *et al.* using *Cmah*<sup>-/-</sup> mice [148].



**Figure 5 Humans have a deletion in the gene encoding CMAH.** Due to the inactive CMAH, human cells only produce Neu5Ac. Most other mammals, including non-human hominids like gorillas or chimpanzees, convert Neu5Ac into Neu5Gc. Both molecules differ in a single oxygen atom (framed).

### ***Role of glycosylation in the host***

The majority of glycans are indispensable in biological systems and their roles are very diverse. Therefore, this chapter only highlights some of the most important features.

Glycans define the physical characteristics of many structures, like plant cell walls or exoskeletons of insects [149, 150]. Moreover, glycosylation is required for correct folding of glycoproteins [151], and to protect them from cleavage by proteases [152]. Sias are fundamentally important for embryonic development, as the disruption of their biosynthesis is lethal in a mouse model [153]. Negatively charged Sias on the cell surfaces lead to the electrostatic repulsion and thus prevent non-specific interactions of cells, e.g. of red blood cells in circulation [154]. In addition, they are important for cell-cell interactions, like the binding of P-selectins (Calcium-dependent lectins) on endothelial cells and glycoproteins on

the surface of neutrophils, which enables neutrophil adhesion to endothelial cells and their “rolling” as part of inflammatory processes [155].

The most important function of glycans, e.g. Sias, is probably to distinguish ‘self’ from ‘non-self’. By covering receptors and antigenic proteins, glycans are recognized as self-associated molecular patterns (SAMPs), which prevent immune responses against host-derived structures [156]. Several members of the family of CD33-related Sia recognizing immunoglobulin superfamily lectins (CD33r Siglecs) have been shown to be involved in the prevention of immune responses. CD33rSiglecs are transmembrane receptors that consist of extracellular sialic-acid binding domains, immunoglobulin domains and often of at least one cytosolic immune-receptor tyrosine-based inhibitory motif (ITIM), which inhibit cell activation and cytokine signalling when the ligand is bound [157].

In contrast, asialoglycoproteins, desialylated structures, are recognized by the hepatic Ashwell-Morell receptor, which removes them from circulation [158]. This is of importance in the clearance of e.g. senescent platelets. These blood components take part in coagulation and regulate their own elimination with the help of neuraminidases, which desialylate their surface-located glycan structures in order to evoke their clearance [159].

### ***Role of glycans during infection***

Many pathogens enter the host via inhalation into the respiratory tract and encounter epithelial surfaces, which are usually covered by a thick and viscous layer, consisting of high molecular weight glycoproteins with many clustered O-linked glycans, the so-called mucins, water, salts, lipids and other highly glycosylated proteins [160, 161]. The mucus traps microbes and particles, which are cleared by cilia movement or coughing [162]. Thus the mucus presents a physical barrier to inhibit invading pathogens.

However, infections and inflammations have been shown to alter glycosylation of the host. One example is the sialylated LewisX blood group antigen. It plays an important role during inflammation [163], as it is upregulated in gastric tissues during *Helicobacter pylori* infections [164], as well as in the respiratory mucins of cystic fibrosis patients [165].

Microbes interact with glycans on the surface of host cells, which promotes pathogenesis [166]. During pneumococcal colonization, adhesion to glycosaminoconjugates, like heparan sulfate, presents one of the first steps in the interaction with epithelial cells [167]. Furthermore, pneumococci engage with the host glycocalyx and produce enzymes and transporters to scavenge and take up Sias (see chapter 1.3.2), but also other sugars, like Gal and GlcNAc, of the glycan strand [168, 169]. In pneumococci, Sia has been shown to promote biofilm formation, nasopharyngeal colonization and bacterial outgrowth from the upper to the lower respiratory tract [170-172]. While intranasal administration of Neu5Ac



enhanced pneumococcal colonization and spread to the lungs and brain of mice, Neu5Gc did not have any effect on pneumococcal carriage or disease [170, 173].

Viruses, like IAV, also interact with Sias on host cells in order to facilitate their entry [35]. Human IAV adheres preferentially to  $\alpha$ 2-6-linked Sias [174], which are present on ciliated cells in the respiratory tract [175].

Interestingly, bacteria have developed strategies to evade the host's immune response by mimicking host glycans or by decorating themselves with Sias. While some bacteria are able to synthesize Sia on their own (e.g. *Escherichia coli* K1 or *N. meningitidis*), others, like *N. gonorrhoeae*, scavenge Sias from host substances and sialylate their surfaces with the help of sialyltransferases [129].

The capsule of group B streptococci (GBS) is shown to contain Sias and represents a major virulence factor [176]. By binding to the CD33r Siglec-9 on neutrophils, the capsular Sia of GBS downregulates its own killing [177]. In contrast, the pneumococcal capsule does not contain Sias, but pneumococci express sialidases, which cleave terminal Sias from the glycan strand [178]. The pneumococcal sialidase NanA has been associated with an unmasking of the CD33r Siglec-5 and subsequent induction of an immune response in macrophages [179]. During sepsis, NanA removes terminal Sias from glycoconjugates of platelets, promoting intravascular coagulation. The Ashwell receptor clears desialylated platelets from the circulation and therefore improves the outcome of this complication [180].

As discussed earlier, PRRs, e.g. TLRs or NLRs recognize PAMPs. Such ligands are often glycan-containing structures, e.g. bacterial LPS or peptidoglycan, or nucleic acids, like bacterial and viral DNA or RNA [181], which play a significant role in host-microbe interactions.

### ***Sialic acid catabolism in bacteria***

Since glucose is not detectable in the respiratory tract of healthy individuals [182], pulmonary pathogens may utilize Sia in order to gain energy in form of nitrogen or carbon [183]. Studies in *E. coli* have shown that Neu5Ac is transported into the bacterial cell and catabolized, leading to the generation of pyruvate and *N*-Acetylmannoseamine (ManNAc), which is, via several steps, converted to Fructose-6-Phosphate, which is part of the glycolysis [184, 185]. Moreover, it was shown that *E. coli* is also able to grow on Neu5Gc as sole carbon source by using the same enzymes as for the catabolism of Neu5Ac [186].

*S. pneumoniae* was predicted to possess all genes required for Sia metabolism [187, 188]. However, in contrast to TIGR4, D39 is not able to ferment Sia due to a frameshift in the gene for the neuraminate lyase [168].

Other serotypes, like 19F and 23F, are shown to be able to grow on Sia Neu5Ac as sole carbon source. As compared to the presence of glucose, growth on ManNAc or Neu5Ac only leads to a much prolonged generation time [168, 172, 189].

### **1.3.2 Pneumococcal virulence factors**

#### ***Capsule***

Pneumococci are surrounded by a thick polysaccharide capsule, which presents a major virulence factor and is the target for current vaccines on the market [4]. Non-encapsulated isolates seldom cause IPD in humans [190] and are significantly attenuated in their virulence using mouse models [191]. The composition of the capsule varies greatly between serotypes [4], and affects the invasive disease potential of the bacteria, although other bacterial factors are also involved [192]. Since most capsular types are negatively charged, electrostatic repulsion prevents pneumococci from being trapped in the mucus and from phagocytosis by immune cells [83]. Moreover, the capsule shields surface proteins and protects pneumococci from recognition and opsonisation by immune cells [193].

Pneumococci of the same serotype can show substantial variations in their capsular thickness, and are therefore divided into opaque and transparent variants. The switch in capsule production, also known as phase variation, allows pneumococci to adapt to different body sites, like the mucosa of the respiratory tract or the blood stream. In contact with epithelial cells, pneumococci produce lower amounts of capsular polysaccharide in order to expose adhesive structures on their surface [194]. These transparent variants were shown to establish a robust colonization in an infant-rat model [195], but hardly caused any sepsis in a mouse model after intraperitoneal challenge. In contrast, all mice succumbed to an intraperitoneal infection with opaque isolates, which produce large amounts of capsule and reduce the detection by the immune system [196].

#### ***Virulence factors involved in sialic acid scavenging and uptake***

*S. pneumoniae* is able to use Sia as carbon source [172, 197]. Genes for Sia removal, uptake and metabolism are encoded in two loci, the *nanAB* and *nanC* loci [178, 184]. These genetic regions harbour up to three different sialidases, with NanA, NanB and NanC to be common to 100%, 96% and 51% of all isolates, respectively [198].

The *nanAB* locus is predicted to contain four transcriptional units [178] and catabolite repression elements, which lead to its transcriptional inhibition in the presence of glucose

[199]. In the absence of glucose, genes within this locus are upregulated in response to Neu5Ac [189, 200].

This thesis focuses on NanA, which is a typical sialidase and scavenges Sias from host glycoconjugates [169, 201]. NanA is located on the bacterial surface and linked to the peptidoglycan cell wall via an LPxTG motif in most pneumococci [202]. Interestingly, in *S. pneumoniae* TIGR4 NanA is secreted due to an authentic frameshift [188].

Despite comprising an enzymatic active part, NanA also consists of a carbohydrate-binding domain. Using a model of experimental meningitis, it was demonstrated that the lectin-like domain is of higher importance than the sialidase domain to achieve adhesion to and invasion into brain endothelial cells, as well as to activate an immune response in these cells [203, 204].

NanA is able to remove both  $\alpha$ 2-3- and  $\alpha$ 2-6-linked Sias from the underlying Gal of the glycan strand [178]. By removing terminal Sias from epithelial cells, potential binding sites are unmasked, and pneumococcal adhesion is promoted [205]. Additionally, desialylation of THP-1 monocytes is reported to stimulate an immune response and to induce cytokine secretion [179]. Moreover, it was suggested that desialylation of host factors, e.g. lactoferrin, interferes with their functionality [178], thus promoting pneumococcal immune evasion. *S. pneumoniae* has also been demonstrated to desialylate host structures of other bacteria in the respiratory tract, like *N. meningitidis* and *H. influenzae*, which is proposed to provide a competitive advantage for pneumococci [206].

NanA expression is increased in transparent compared to opaque variants [178], and it was shown to be beneficial during colonization in a chinchilla model [201]. Moreover, *nanA* transcription was strongly increased in bacteria isolated from the nasopharynx, lungs and brain of mice in comparison to blood [207, 208].

Several groups observed that NanA is also required for colonization of the nasopharynx and lungs using a murine model of intranasal challenge [209, 210]. Moreover, a recent study reported that NanA-mediated exposure of Gal *in vivo* in the nasopharynx promotes pneumococcal biofilm formation [211]. During sepsis, NanA is demonstrated to be redundant for pneumococcal survival in the vascular system [210, 212, 213], but there are studies showing a reduced virulence of NanA-deficient mutants in the blood stream [209, 213]. Moreover, NanA has been shown to promote pneumococcal meningitis in murine *in vivo* models [203].

Three different Sia transporters have been predicted in the *nanAB* and *nanC* loci of *S. pneumoniae* TIGR4, a solute symporter SP1328, and two ABC transporters SP1688-90 and SP1681-83 [187]. The latter was identified as the main Sia transporter, and named SatABC. Moreover, it has been shown to play an important role in nasopharyngeal colonization of mice [172].

### ***Two-component system CiaRH***

Two-component systems (TCS) allow bacteria to sense and respond to environmental signals. They comprise a membrane-bound histidine kinase (HK), and a cytoplasmic response regulator (RR). The HK detects the signal on the outside of the cell, is autophosphorylated and phosphorylates the RR, which alters its conformation in order to bind to specific promoter regions and thus regulates transcription [214].

In *S. pneumoniae*, 13 different TCS and a single RR have been annotated, of which the majority affects pneumococcal virulence [215]. During the analysis of mutants resistant to the  $\beta$ -lactam antibiotic cefotaxime, TCS05, also known as CiaRH, was identified and a connection with the pneumococcal cell wall machinery was suggested [216]. Besides sensitivity to cefotaxime, CiaRH was also shown to affect autolysis. Deletion mutants in *ciaR* lyse quickly upon entry into stationary phase or in response to cell wall inhibitors, like vancomycin [217]. Other studies revealed that CiaR regulates the expression of several genes involved in biosynthesis of the pneumococcal cell wall [218, 219].

The use of microarrays and solid-phase DNA binding assays also identified the high-temperature requirement A gene (*htrA*) to be regulated by CiaRH [218, 219] (see below), and the contribution of CiaRH to oxidative stress resistance and virulence in rodents, using nasopharyngeal colonization and pneumonia models, was demonstrated [215, 219, 220]. Moreover, CiaRH was shown to play a role in bacteriocin production [218].

A mutation in the HK CiaH that mimicked its activation, led to the complete abolishment of competence [216], while a mutation in RR CiaR restored competence [221]. Another study reported the upregulation of the competence operon in a *ciaRH*-mutant [219]. However, a direct interaction between CiaR and genes involved in competence could not be observed [218]. Moreover, CiaR expression is also regulated by competence, as *ciaR* transcription is increased in the late phase of competence, which was suggested to promote re-entry into the non-competent state [222].

The external stimulus for the activation of CiaH is not known yet, but it might be induced upon stress. Alignments of promoter sequences targeted by CiaR identified its binding motif (NTTAAG-N5-TTTAAG) [223]. Moreover, by cloning CiaR-regulated promoter regions in front of a  $\beta$ -galactosidase gene (*lacZ*), it was shown that the position of the binding motif affects transcription control. If the motif is located upstream of the transcriptional start site, CiaR will positively regulate gene transcription. In contrast, if the binding site is located on the other DNA strand inside the transcribed region, CiaR downregulates promoter activity [224].

Bioinformatic analysis identified five *cia*-regulated small RNAs (csRNA). Their functional analysis implicated a role in autolysis, although the phenotype that was caused by their

deletion was not as pronounced as in a *ciaR* null mutant [224]. Moreover, a function in competence [225, 226] and virulence in mice after intranasal challenge was suggested [227].

CiaRH is conserved in streptococci but absent in other bacterial species [223]. In order to bind to promoter fragments and to regulate transcription, CiaR needs to be phosphorylated. Interestingly, depending on the growth media used, CiaH is or is not required to activate CiaR [221, 228]. Intracellular acetyl phosphate (AcPh) production is strongly regulated by the streptococcal pyruvate oxidase (SpxB) (see below) [229]. Deletion of SpxB leads to a significant downregulation of CiaR-regulated genes, suggesting that intracellular AcPh is capable of phosphorylating this RR [230].

### ***Streptococcal pyruvate oxidase (SpxB)***

The pneumococcal pyruvate oxidase SpxB is located in the cytoplasm and catabolizes the decarboxylation of pyruvate leading to the production of AcPh and H<sub>2</sub>O<sub>2</sub> [231]. Pneumococci generate high concentrations of H<sub>2</sub>O<sub>2</sub>, which enables them to outcompete other respiratory pathogens, like *N. meningitidis* or *H. influenzae*, present in the same niche [232]. Interestingly, pneumococci themselves lack catalase [233]. Instead HtrA, as mentioned above, was shown to promote resistance to oxidative stress [234].

SpxB has been associated with the pneumococcal capability to bind to glycoconjugates, which are found on endothelial cells and lung epithelial cells [231]. A mutation in *spxB* leads to drastically reduced AcPh and acetyl-CoA amounts in the bacteria, which are suggested to mediate adhesin expression. On one hand, acetyl-CoA is required for the biosynthesis of fatty acids, amino acids and the cell wall, and subsequently affects fundamental structures of the bacterial cell [231]. On the other hand, AcPh is known to act as a phospho-donor for bacterial TCS [235]. A deletion of *spxB* downregulates *ciaR* mRNA levels and thus CiaR-regulated genes [230], which might affect adhesin expression.

H<sub>2</sub>O<sub>2</sub> has been shown to inhibit ciliary beating and cell integrity of human nasal epithelial cells [236]. It was also reported to induce host cell apoptosis in meningitis models, using microglia cells and rat primary neurons [237], and in lung epithelial cells by introducing DNA double-strand breaks [238]. It has been reported to evoke stress responses in respiratory epithelial cells and the secretion of inflammatory mediators [239]. Moreover, a deletion of *spxB* in serotype 4 pneumococci reduces the production of capsular polysaccharides [240]. In summary, SpxB is a main pneumococcal virulence factor and its importance has been demonstrated using several *in vivo* models, e.g. of colonization, pneumonia or bacteraemia [231, 240].

### ***High temperature requirement A (HtrA)***

HtrA is a serine protease that is highly conserved among bacteria and located on the pneumococcal surface [241]. It plays an important role in protein quality control, as it is described to have chaperone and protease functions. HtrA promotes protein folding at lower temperatures, and fosters degradation of misfolded proteins at higher temperatures [242].

In pneumococci, HtrA promotes growth at higher temperatures [234] and its expression is regulated by the pneumococcal TCS CiaRH [218, 219]. Moreover, HtrA mediates pneumococcal resistance to oxidative stress [234], which is caused by ROS-producing immune cells like macrophages or neutrophils [243, 244].

Several reports have highlighted an effect on competence by HtrA, although they presented opposing results. Some studies have shown that HtrA contributes to the inhibitory effect of CiaRH on competence by degrading competence-stimulating peptides [241, 245], while others have demonstrated a decrease in competence in a mutant lacking HtrA [234].

HtrA of *H. pylori* and *Campylobacter jejuni* have been shown to degrade E-cadherin, which leads to the disruption of tight junctions, promoting bacterial invasion [246, 247]. In *S. pneumoniae*, HtrA has also been associated with virulence. Mutants in *htrA* showed lower colonization rates and led to lower bacterial counts in the lungs of rodents after an intranasal challenge, as well as a reduced rate of bacteraemia after intravenous infection [219, 234, 248].

### ***Pneumolysin***

Pneumococci were the first bacteria, which were demonstrated to lyse red blood cells [249]. This lysis is caused by pneumolysin, a major virulence factor, which is expressed by nearly all pneumococcal isolates [250]. It is a 52 kDa binding monomer that forms a ring of 30-50 subunits, thus interfering with membrane integrity and leading to cell lysis [251, 252].

Pneumolysin binds to cell membranes via cholesterol [251], but has also been reported to adhere to glycan structures. It was shown to attach to mannose and sialylated LewisX blood group antigens on the cell surface, since the haemolytic activity of pneumolysin was strongly inhibited after pre-incubation with these sugars [253, 254]. Sialyl LewisX is found on various immune cells like neutrophils or monocytes [255, 256] and has been shown to play a significant role in the first step of extravasation of these cells [163].

To interact with host cells, pneumolysin has to be present on the outside of bacteria. Interestingly, pneumolysin was found to be located in the cytoplasm [257] and does not contain a signal sequence, which would direct its translocation to the extracellular space [258]. Yet, it is detected non-covalently attached to the bacterial cell wall [259]. For many years, pneumolysin was thought to be released during pneumococcal autolysis. However,

recent studies demonstrated an autolysis-independent secretion of the cytotoxin, suggesting an active transport of pneumolysin out of the bacterial cell [260-262].

Pneumolysin activates the immune system. It promotes bacterial adhesion to epithelial cells [263] and reduces ciliary movement in the lungs [236]. At sublytical concentrations, pneumolysin stimulates the secretion of cytokines, like IL-6, TNF $\alpha$  or IL-1 $\beta$ , by different cell types, e.g. DCs and monocytes [71, 264]. However, other reports showed an inhibition of inflammatory responses by pneumolysin [265, 266]. While some studies demonstrated that host cells sense pneumolysin in a TLR4-mediated way, others observed TLR4-independent but inflammasome dependent signalling in response to this cytotoxin [71, 267]. Its importance *in vivo* was demonstrated using mouse models of pneumonia and bacteraemia, as a mutant deficient in pneumolysin was severely attenuated in virulence [250, 268]. Moreover, pneumolysin-mediated inflammation was shown to endorse pneumococcal shedding and its transmission between hosts [269].

### ***Pathogenesis of coinfections with influenza and pneumococci***

Preceding influenza infections significantly worsen the outcome of a secondary pneumococcal infection. The pathogenesis of coinfections with influenza and pneumococci is regulated by an interplay of viral, bacterial and host factors.

Human influenza viruses bind to  $\alpha$ 2-6-linked Sias [174], which are found in the respiratory tract of men [175]. The infection with the virus induces drastic changes in the respiratory tract, which promote pneumococcal infections. Influenza infections foster secondary bacterial colonization by damaging the respiratory epithelium, which prevents ciliary clearance [270]. Moreover, viral sialidases have been shown to cleave Sias from epithelial surfaces and mucins, thereby exposing host receptors, enabling increased bacterial adhesion [271, 272], and providing an energy source for pneumococcal growth and translocation from the upper respiratory tract into the lungs [171].

At early stages of influenza infections, the number of alveolar macrophages in the lungs are severely decreased [273]. It takes up to 2 weeks until the macrophage reservoir is restored by the infiltration of new macrophages. Thus, the host susceptibility to secondary bacterial infections is increased during this period [35]. Furthermore, influenza infections increase the production of IFN- $\gamma$ , which is shown to inhibit macrophage function by downregulating MARCO, a scavenger receptor known to promote pneumococcal phagocytosis [274]. Following influenza infections, an inhibition of ROS-mediated bacterial killing by macrophages was also demonstrated [275].

Although macrophage numbers are decreased, neutrophil numbers are increased at day 7 post influenza infection. However, their role in bacterial influenza coinfections is still debated. While some studies report that increased inflammatory processes due to neutrophil influx

correlate with higher bacterial burden and lower survival rates in coinfecting versus single infected mice, others described a redundancy of neutrophils in coinfection settings, as their depletion did not affect the disease outcome [276].



## **2 AIMS**

Since glycans are highly abundant in nature, they play an important role in host-pathogen interactions. Especially Sias, the terminal sugars of the glycan strand, serve as receptor for many pathogens. The main scope of this thesis was to determine virulence regulation in the human-adapted pathogen *S. pneumoniae* in response to Sias, in particular Neu5Ac and Neu5Gc, which are mainly abundant in humans and other mammals, respectively. Despite focusing on infections with pneumococci only, preceding influenza infections, which are known to lead to severe complications, were also taken into account. Moreover, the activation of host immune responses by pneumococcal glycan structures was investigated.

### **2.1 SPECIFIC AIMS**

#### **Paper I**

To study pneumococcal pathogenesis and the expression of virulence factors in response to the Sias Neu5Ac and Neu5Gc, as well as to determine how these two sugars effect intracellular signalling in pneumococci.

#### **Paper II**

To analyse pneumococcal interactions with Neu5Ac or Neu5Gc on host cells and to characterise the effect of pneumococcal toxins on the evoked host response *in vitro* and *in vivo*.

#### **Paper III**

To explore alterations of the host immune response in the lower respiratory tract after influenza infection and their consequences on coinfection with *S. pneumoniae*, as well as to investigate pneumococcal adaptations to this environment.

#### **Paper IV**

To identify pneumococcal structures that activate immune responses by dendritic cells and to find a possible mechanism for the enhanced cytokine secretion, observed in influenza/pneumococcal coinfections.



### 3 METHODOLOGICAL CONSIDERATIONS

This chapter gives a general overview of infectious agents and methods that have been used for the work of this thesis. For a more detailed description of the protocols, see Material and Methods sections of the respective papers.

#### 3.1 BACTERIAL AND VIRAL STRAINS AND THEIR GROWTH CONDITIONS

*S. pneumoniae* was cultured in C medium with yeast extract (C+Y) [277]. In some experimental set-ups, monosaccharide-free C+Y was supplemented with either 12 mM glucose, Neu5Ac or Neu5Gc. Liquid cultures or bacteria on plates were grown at 37°C in a water bath or at 37°C and 5% CO<sub>2</sub> in an incubator, respectively.

Throughout all the studies in this thesis, encapsulated and invasive pneumococcal serotype type 4, TIGR4 (ATCC BAA-334) [188] or its non-encapsulated derivative T4R (paper II and IV) [278] were used. *In vitro* challenges with T4R were shown to lead to similar numbers of phagocytosed bacteria and to comparable amounts of cytokines produced by the cells compared to studies using an opsonized TIGR4 [266].

Bacterial mutants were constructed by overlap polymerase chain reaction (PCR) of the upstream region, an erythromycin cassette and the downstream region of the target gene [279]. The PCR product containing these three DNA-fragments was then transformed into *S. pneumoniae* TIGR4 or T4R, which was pre-incubated with the competence-stimulating peptide CSP-2 in order to induce competence. All mutations were confirmed by PCR and sequencing.

Since coinfections of IAV and pneumococci are a major cause of mortality, paper III and IV studied different aspects of preceding influenza infections followed by pneumococcal challenges. In paper III, we used the mouse-adapted influenza virus strain PR8/A/34, while in paper IV we used the human-adapted X31. Both were propagated on Madin-Darby canine kidney (MDCK) cells.

## 3.2 CHARACTERIZATION OF PNEUMOCOCCAL PHENOTYPES

### 3.2.1 Sialidase activity

Pneumococcal sialidases are major virulence factors [209, 210]. In order to measure sialidase activity of bacterial lysates in paper I, a protocol by Manco *et al.* was used [209]. 2-O-(p-nitrophenyl)-D-N-acetylneuraminic acid was added to the samples, followed by an incubation at 37°C. Sialidases are able to cleave this substrate into Neu5Ac acid and p-nitrophenol, which has a yellow color. Thus, the absorbance at 405 nm was measured. The values were compared to a standard curve with known concentrations of the sialidase of *Arthrobacter ureafaciens* and normalized to total protein content in order to determine sialidase activity of the sample.

### 3.2.2 Production of hydrogen peroxide and acetyl phosphate

The pyruvate oxidase (SpxB) is a pneumococcal virulence factors and responsible for the production of H<sub>2</sub>O<sub>2</sub> [231]. To determine amounts of H<sub>2</sub>O<sub>2</sub> in samples of paper I and III, we used a previously described method [229]. 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid) and horseradish peroxidase were added onto the sample. After incubation at room temperature, the absorbance at 560 nm was measured. Total amounts of H<sub>2</sub>O<sub>2</sub> were calculated with the help of a standard curve, which was produced with known concentrations of H<sub>2</sub>O<sub>2</sub> and normalized to bacterial numbers in the samples.

SpxB was reported to affect intracellular acetyl phosphate (AcPh) levels [231]. In paper I, AcPh content inside the bacterial cell was studied according to the protocol by Pruss and Wolfe [280]. In order to measure total AcPh amounts, it was used to convert adenosine diphosphate (ADP) to adenosine triphosphate (ATP) by an acetate kinase, and quantified using a bioluminescence kit. Finally, the results were compared to a standard curve with known AcPh amounts and normalized to total protein in the sample.

### 3.2.3 Survival in response to hydrogen peroxide

Although pneumococci have been shown to be catalase-negative, they developed mechanisms to resist ROS [84, 234]. To study the sensitivity of pneumococci to H<sub>2</sub>O<sub>2</sub> in paper I and III, the method published by Ibrahim *et al.* was used [220]. Bacteria were incubated in the presence of H<sub>2</sub>O<sub>2</sub> at 37°C. The percentage of surviving bacteria was calculated by dividing viable bacterial numbers after exposure by viable bacterial numbers before exposure to H<sub>2</sub>O<sub>2</sub>.

### 3.3 IN VITRO AND EX VIVO MODELS

#### 3.3.1 Cells

Since *S. pneumoniae* is a human-adapted pathogen [281], we performed most of the *in vitro* experiments with human cell lines or compared cells with either Neu5Ac- or Neu5Gc-containing glycoconjugates.

In paper II, human cell lines THP-1 (ATCC TIB-202) and A549 (ATCC CCL-185) derived from acute monocytic leukemia or epithelial lung cell carcinoma, respectively, were used. To differentiate THP-1 cells into macrophage-like cells, they were incubated with phorbol myristate acetate. Cell lines present a good *in vitro* model in order to mimic the *in vivo* setting. However, the immortalization of the cells can also lead to differences in their phenotype, e.g. receptor expression levels. Therefore, we additionally performed experiments with primary alveolar macrophages, which were isolated from the BAL from uninfected C57BL/6 wild-type (wt) or *Cmah*<sup>-/-</sup> mice. While wt mice mainly present Neu5Gc on the surface of their cells, *Cmah*<sup>-/-</sup> mice have a human like deletion in exon 6 of the *Cmah* gene [282], leading to the lack of Neu5Gc.

In paper IV, primary monocytes from buffy coats of healthy human donors were isolated and cultured with granulocyte-macrophage colony-stimulating factor (GM-CSF) and IL-4 to differentiate them into DCs. Successful differentiation was confirmed by the presentation of the surface markers CD1a and CD11c.

Human embryonic kidney 293 cells (HEK293, ATCC CRL-1573) were used in paper IV. These cells were stably transfected with plasmids either expressing TLR3 or TLR4 and luciferase.

#### 3.3.2 Sialic acid feeding of cells

The Sia profiles differ greatly between humans and most other mammals [133, 134]. To study the effect of Neu5Ac- and Neu5Gc-presentation by host cells on pneumococcal virulence in paper II, a previously published method was used [283, 284]. We supplemented the RPMI 1640 cell medium with Neu5Ac and heat-inactivated human AB serum or Neu5Gc and heat-inactivated fetal bovine serum. The Neu5Ac- and Neu5Gc-presentation by the cells was confirmed by staining with *Sambucus nigra* lectin (SNA), *Maackia amurensis* lectin II (MAA II) and an antibody raised against Neu5Gc.

### 3.3.3 Measurement of cytokine response and cell toxicity

In all papers, pro-inflammatory cytokines, like human or murine TNF $\alpha$  or interleukin 6 (IL-6), IL-12, and chemokines, like human IL-8 or murine KC or macrophage inflammatory protein 2 (MIP-2), were determined in cell supernatants or BAL of mice using enzyme-linked immunosorbent assay (ELISA).

## 3.4 IN VIVO MODELS

Pneumococci have humans as their natural host [281] but different animal models, like rats and mice, have been described as a suitable model to study pneumococcal infections [285]. *In vivo* models allow studying pneumococcal-host interactions in a complex environment, which also takes the immune system as a whole into account. In paper I, II and III, wt and/or *Cmah*<sup>-/-</sup> mice have been used. Besides, diminishing the synthesis of Neu5Gc, an inactivation of *Cmah* in the mice caused age-related hearing loss and defects in wound healing without affecting the recruitment of immune cells [282]. Similar to humans, *Cmah*<sup>-/-</sup> mice are also demonstrated to exhibit an over-reacting immune system [143, 144]. A recently reported *dock2* mutation in some *Cmah*<sup>-/-</sup> mice, which occurred due to backcrossing into commercially available C57BL/6 mice [286], was not present in the mice used in this thesis.

In paper I and II, mice were infected intranasally with 20  $\mu$ l PBS containing *S. pneumoniae*, presenting the natural route of infections by pneumococci. About half of the inoculum reached the lungs directly during infection, allowing us to study the development of pneumonia and the progress into septicaemia in the mice. In paper I, we also infected the mice intravenously, in order to study pneumococcal infections in the blood stream. In paper III, we challenged the mice intranasally with influenza. To study pneumococcal/influenza coinfections, pneumococci were instilled intratracheally, which allowed us to deliver the whole bacterial inoculum into the lower respiratory tract.

## 3.5 ETHICAL CONSIDERATIONS

All experiments were performed in accordance with the local ethical committee (Stockholms Norra djurförsöksetiska nämnd). Mice were kept with food and water *ad libitum* and 12 hours light/dark cycles. The health status of the mice was controlled regularly.

## 4 RESULTS AND DISCUSSION

### 4.1 PAPER I

#### ***Streptococcus pneumoniae* senses a human-like sialic acid profile via the response regulator CiaR**

When pulmonary pathogens, like *S. pneumoniae*, enter the respiratory tract, they come in contact with Sias in mucins [166, 287]. Pneumococci are adapted to humans as their natural host [281]. Thus, the first study included in this thesis determined pneumococcal virulence in response to the Sias Neu5Ac and Neu5Gc, which are predominantly found in mucins of humans or of most other mammals, respectively. Moreover, we compared expression levels of pneumococcal virulence factors in response to these sugars and elucidated how Sia catabolism affects intracellular signalling in pneumococci.

To study pneumococcal pathogenesis in response to Neu5Ac and Neu5Gc, we challenged C57BL/6 wt and *Cmah*<sup>-/-</sup> mice with pneumococci. While wt mice are able to convert Neu5Ac to Neu5Gc, *Cmah*<sup>-/-</sup> mice, like humans, overproduce Neu5Ac and are not able to synthesize Neu5Gc [282, 288]. We chose an *in vivo* model of intranasal challenge, since it presents the natural route of infection by pneumococci. At 24 hours post infections, we detected significantly higher bacterial numbers in lungs, BAL and the blood as well as reduced survival rates of *Cmah*<sup>-/-</sup> compared with wt mice. Hence, pneumococci trigger stronger symptoms of disease in *Cmah*<sup>-/-</sup> mice in comparison to wt mice. In agreement with our results Trappetti and colleagues observed higher bacterial counts in the lungs of mice after intranasal administration of Neu5Ac but not Neu5Gc [170].

Besides being able to utilize Neu5Ac as carbon source [172, 197], our data indicate that *S. pneumoniae* can also grow on Neu5Gc. Since pneumococci are human-adapted pathogens, we propose that pneumococcal virulence factors mediate increased virulence in response to human-like Sia profiles compared to Neu5Gc. We decided to study the activity of sialidases, since pneumococcal sialidase NanA was shown to bind to Sia and to remove it from the glycan strand [178]. Moreover, sialidases of other bacteria, viruses and mammals have been reported to preferentially adhere to and cleave Neu5Ac- compared with Neu5Gc-containing substrates [289-291]. To catabolize Sia, it needs to be taken up into the bacterial cell [172] and both NanA and the main Sia transporter SatABC have recently been demonstrated to be upregulated in response to Neu5Ac [189, 200]. We isolated pneumococcal RNA from *in vitro* conditions using monosaccharide-free C+Y medium supplemented with either Neu5Ac or Neu5Gc, and *in vivo* from the nasopharynx of wt and *Cmah*<sup>-/-</sup> mice. *NanA* transcription, as well as sialidase activity was increased in Neu5Ac-incubated bacteria compared to Neu5Gc-treated ones. Moreover, bacteria isolated from the nasopharyngeal lavage of *Cmah*<sup>-/-</sup> mice had higher *nanA* mRNA levels in comparison to those isolated from wt mice. Additionally, the substrate binding protein SatA of the main Sia transporter was also higher transcribed in

response to Neu5Ac as compared with Neu5Gc. The differential transcription of *nanA* in TIGR4 in response to the two sugars was dependent on the three known Sia transporters, SatABC, SP1688-90, and SP1328 [184, 187], which suggest that pneumococci upregulate virulence factors in response to Neu5Ac compared with Neu5Gc after it has been taken up into the bacterial cell.

To test whether pneumococcal TCS are involved in the Sia-mediated signalling, we monitored transcription rates of all known pneumococcal RR, as well as growth rates of their deletion mutants on Sia and glucose. Our data reveal that a mutant deficient in CiaR had severe growth defects on Neu5Ac and Neu5Gc but not on glucose. Moreover, mRNA expression of RR05, also known as CiaR, was increased in bacteria treated with Neu5Ac compared with Neu5Gc but only if the Sia transporters were present. Hence, we hypothesized that this RR could be involved in intracellular signalling in response to Sia or its metabolic compounds. The TCS CiaRH is known to control the expression of HtrA, which mediates pneumococcal resistance to oxidative stress [234]. Consequently, we measured *htrA* transcription and pneumococcal survival in response to H<sub>2</sub>O<sub>2</sub> and found both to be increased after incubation in Neu5Ac compared with Neu5Gc.

CiaR needs to be phosphorylated in order to bind to target promoter sequences [228]. Interestingly, it was shown that its phosphorylation does not depend on the HK CiaH, instead intracellular AcPh can also serve as phosphor-donor [230]. In pneumococci, the pyruvate oxidase SpxB is the main producer of AcPh and H<sub>2</sub>O<sub>2</sub> [229, 231]. Moreover, its expression was shown to affect CiaR transcription [230]. After treatment with Neu5Ac, we observed higher *spxB* transcription as well as increased H<sub>2</sub>O<sub>2</sub> and AcPh amounts inside pneumococci in comparison to Neu5Gc treatment. Noteworthy, an *spxB*-deficient mutant did not only downregulate *ciaR* transcription, it also decreased *nanA* levels significantly.

CiaR, NanA as well as SatABC are well-known pneumococcal virulence factors [172, 209, 210, 215, 292]. While intranasal infections with wt TIGR4 resulted in higher bacterial numbers in the lungs and blood of *Cmah*<sup>-/-</sup> compared with wt mice, challenges with TIGR4Δ*ciaR*, TIGR4Δ*nanA* or TIGR4Δ*satABC* led to comparable cfu in the lungs of wt and *Cmah*<sup>-/-</sup> mice without invading into the bloodstream. Thus, CiaR, NanA and SatABC contribute to a more severe outcome of pneumococcal disease in *Cmah*<sup>-/-</sup> compared with wt mice.

In TIGR4, the genes required for Sia retrieval and metabolism are encoded in the *nanAB* and the *nanC* loci [178, 184]. The *nanAB* locus harbours catabolite repression elements [189, 200], which suppress the transcription of genes inside the locus in the presence of glucose. Also, CiaR is reported to be downregulated in response to glucose [293]. Since we did not observe any differences in bacterial numbers and survival rates between wt and *Cmah*<sup>-/-</sup> mice after intravenous challenge, we propose that NanA, SatABC and CiaR facilitate a faster disease progression in the glucose-free respiratory tract [294] of *Cmah*<sup>-/-</sup> compared with wt mice, but not in the blood stream with high glucose concentrations [295].



This study demonstrates a possible mechanism for increased pneumococcal virulence in response to Neu5Ac in contrast to Neu5Gc. Our data indicate that pneumococci are able to signal in response to Sias via the RR CiaR. We propose a model, in which the presence of Neu5Ac increases expression of sialidase NanA and Sia transporter SatABC, leading to increased scavenging of Sias and higher amounts of Sia or its metabolic products, e.g. pyruvate, inside the bacterial cell. Subsequently, SpxB expression is upregulated, leading to enhanced levels of AcPh and H<sub>2</sub>O<sub>2</sub>. Increased AcPh amounts inside the bacterial cell upregulate CiaR, which mediates increased resistance to ROS via HtrA and upregulates NanA expression.

## 4.2 PAPER II

### Human sialic acid profiles mediate increased pneumococcal adhesion and immune evasion

The second study of this thesis further characterised pneumococcal host tropism. Since we observed an upregulation of pneumococcal virulence factors in response to Neu5Ac as compared with Neu5Gc in paper I, we aimed at investigating how pneumococci interact with Neu5Ac- or Neu5Gc-presenting host cells and how this interplay affects the host's immune response.

Since it was shown that virulence factors of several human-adapted pathogens favourably bind to human-like sialic acid profiles [140, 142], we set up an *in vitro* model to study pneumococcal interactions with cells presenting either Neu5Ac or Neu5Gc. Sia feeding has been reported to modify glycan structures of several cell lines [283, 284]. We used macrophage-like THP-1 cells and a lung epithelial cell line (A549). By supplementing the cell culture media with Neu5Ac and human serum or Neu5Gc and fetal calf serum, we could detect  $\alpha$ 2-3- and  $\alpha$ 2-6-linked Sias by staining the cells with the lectins MAAII and SNA, respectively. However, only after Neu5Gc supplementation could Neu5Gc be detected on the surface of both cell lines using anti-Neu5Gc antibodies. In contrast, cells that were cultured in the presence of Neu5Ac were not recognized by the anti-Neu5Gc antibody. We propose that they most likely present Neu5Ac, although this remains to be experimentally confirmed.

Next, we studied whether Neu5Ac- or Neu5Gc-supplementation of the growth media affects pneumococcal adhesion to cells. We used a non-encapsulated variant of TIGR4, T4R, since it was shown to behave similar to opsonized wt TIGR4 in *in vitro* phagocytosis assays [266]. By performing cell adhesion and uptake assays, we detected higher numbers of cell-associated bacteria when cells were cultured in the presence of Neu5Ac as compared with Neu5Gc. However, we excluded the pneumococcal capsule as determinant of this phenotype, as both wt TIGR4 and T4R adhered better to Neu5Ac- than to Neu5Gc-incubated cells. To test whether increased pneumococcal adhesion to Neu5Ac as compared with Neu5Gc-presenting cells, can be extrapolated to primary cells, we isolated primary alveolar macrophages from uninfected wt and *Cmah*<sup>-/-</sup> mice. After challenging the cells with pneumococci, we observed higher adhesion rates to macrophages from *Cmah*<sup>-/-</sup> compared with those from wt mice.

The pneumococcal pyruvate oxidase SpxB and pneumolysin are potent virulence factors and stimulate inflammatory processes [236]. In line with observations of other groups, we detected lower adhesion rates of *spxB* and pneumolysin deficient mutants compared with wt bacteria [263, 296]. Since inflammatory processes are reported to alter the glycan structures on host cells [164, 165], we proposed that these pneumococcal cytotoxins could upregulate glycosylations on cells and hence provide additional attachment sites for bacteria. Moreover, mutants in *spxB* produce significantly lower AcPh amounts than wt bacteria [297]. AcPh is

shown to be able to phosphorylate RR CiaR, a major pneumococcal regulator [230], thus expression levels of adhesins might also be reduced. It will be interesting to test which pneumococcal adhesins favourably bind to Neu5Ac than to Neu5Gc.

More bacteria also lead to higher levels of pneumococcal virulence factors, like SpxB or pneumolysin. Recently, it was shown that the presence of SpxB and pneumolysin cause host cell apoptosis [210, 237, 250, 297, 298]. Consequently, the increased bacterial burden on Neu5Ac-incubated cells might cause higher cytotoxicity in comparison to lower bacterial numbers associated with Neu5Gc-fed cells.

Like T4R, mutants in *spxB* and pneumolysin adhered better to Neu5Ac- than to Neu5Gc-cultured cells. T4R and T4R $\Delta$ *spxB* were also taken up more into Neu5Ac- compared to Neu5Gc-treated cells. In contrast, there was no significant difference in phagocytosis of T4R $\Delta$ *ply* between Neu5Ac- and Neu5Gc-fed cells. Since pneumolysin has been demonstrated to bind to sialylated carbohydrates on cells [254], we speculate that extracellular pneumolysin facilitates higher pneumococcal adhesion to Neu5Ac as compared with Neu5Gc on host cells.

To study the host's immune response, we measured the production of IL-8 after pneumococcal infection of Neu5Ac or Neu5Gc-incubated THP-1 cells. IL-8 concentrations were reduced in cells fed with Neu5Ac compared with Neu5Gc after challenge with T4R and T4R $\Delta$ *spxB* but not after infection with T4R $\Delta$ *ply*. It is tempting to speculate that lower IL-8 levels are caused by higher cytotoxicity in Neu5Ac-fed cells, but this remains to be tested.

To determine the importance of Neu5Ac- and Neu5Gc-presenting cells in pneumococcal infections *in vivo*, we challenged wt and *Cmah*<sup>-/-</sup> mice intranasally with TIGR4, TIGR4 $\Delta$ *spxB* or TIGR4 $\Delta$ *ply*. We observed similar bacterial numbers in BAL of wt and *Cmah*<sup>-/-</sup> mice after challenge with TIGR4 or its isogenic mutants in *spxB* or pneumolysin. However, we detected lower total cell numbers, fewer neutrophils as well as lower MIP-2 and TNF $\alpha$  concentrations in BAL of *Cmah*<sup>-/-</sup> compared with wt mice after challenge with TIGR4 but not after infection with TIGR4 $\Delta$ *spxB* or TIGR4 $\Delta$ *ply*. KC concentrations were lower in *Cmah*<sup>-/-</sup> than in wt mice after infection with TIGR4 and TIGR4 $\Delta$ *spxB*. Infections with TIGR4 $\Delta$ *spxB* or TIGR4 $\Delta$ *ply* almost did not cause any neutrophil influx in the BAL. To exclude an immunodeficiency of *Cmah*<sup>-/-</sup> mice, we administered LPS intranasally into wt and *Cmah*<sup>-/-</sup> mice. Both mice showed a similar immune response and recruited comparable numbers of cells into the airway.

In summary, we observed higher pneumococcal adhesion rates to cells expressing human-like Sia profiles. Moreover, higher bacterial numbers resulted in lower IL-8 secretion, which was mainly mediated by the pneumococcal cytotoxin pneumolysin. A recently published study reported that also *E. coli* adheres better to cells of *Cmah*<sup>-/-</sup> than wt mice. The authors observed increased bacterial killing after infecting macrophages of *Cmah*<sup>-/-</sup> mice with sublethal doses of *E.coli* but a higher susceptibility to endotoxins in the knock-out mice after intraperitoneal challenge. They hypothesized that the CMAH loss provided

an advantage to clear minor infections, while it can be harmful during severe infections [135].

In line with our *in vitro* data, we observed lower cytokine and chemokine concentrations as well as lower numbers of infiltrating neutrophils in the BAL of *Cmah*<sup>-/-</sup> than in wt mice after challenge with TIGR4. Thus, we propose that higher pneumococcal adhesion rates to alveolar epithelial cells and/or macrophages of *Cmah*<sup>-/-</sup> mice compared with wt mice mediate increased cytotoxicity in the knockout mice. Macrophages and epithelial cells produce cytokines and chemokines, which evoke an influx of neutrophils to the site of infection. Hence, higher cell death rates in these cells might lead to lower secretion rates of inflammatory mediators and lower numbers of recruited neutrophils. However, further experiments are required to test this hypothesis.

Additionally, our data indicate that using cell culture medium supplemented with fetal bovine serum in *in vitro* assays might skew the experimental outcome for human-adapted pathogens, as they might preferentially bind to Neu5Ac and not to Neu5Gc.

### 4.3 PAPER III

#### **Mechanism for enhanced bacterial burden in the lower respiratory tract of mice during influenza/pneumococcal coinfection**

The third study of this thesis determined influenza-mediated alterations in the respiratory tract and their consequences for pneumococcal adaptation and virulence regulation.

To set up an *in vivo* model for studying influenza/pneumococcal coinfections, we challenged mice with pneumococci 7 days post influenza infection. At this time point, the mice were recovering from the primary influenza infection. Additionally, humans are also shown to be highly susceptible to bacterial superinfections about one week post influenza infection [299]. In line with earlier publications [274, 300], we detected higher bacterial burden in the BAL of influenza/pneumococcal coinfecting mice compared to those, which were challenged with pneumococci only.

Viral sialidases are known to cleave terminal Sias from the glycan strand [301], hence they increase the concentration of free Sia in the respiratory tract. Siegel and colleagues showed that pneumococci utilize virus-released Sias in the nasopharynx, which promotes pneumococcal colonization and translocation into the lungs [171]. To test whether free Sias also foster pneumococcal dissemination in the lower respiratory tract (LRT), we infected mice with TIGR4 and its isogenic mutant in the main Sia transporter *SatABC*. Like wt TIGR4, infections with a *satABC*-deficient mutant led to increased bacterial numbers in the BAL and lungs of coinfecting compared with single-infected mice. Hence, pneumococcal outgrowth in the LRT is independent on growth on Sia.

To determine total numbers of immune cells and to investigate their contribution to the host response in the LRT, we isolated cells from BAL of untreated, influenza infected, pneumococcal challenged and coinfecting mice. As it was reported by other groups [273], the number of macrophages was decreased in influenza and coinfecting mice, which might contribute to a defect in bacterial clearance [273, 274, 302, 303]. In contrast, cytokine release and neutrophil numbers were increased after challenge with influenza and in coinfecting mice. Thus, although bacterial clearance might be reduced, recruitment of neutrophils and cytokine secretion mechanisms seem to be functional in our model.

AMPs are known to play an importance role in the innate immune response to invading pathogens. To measure the production of AMPs and collectins in alveolar macrophages, neutrophils and epithelial cells, we isolated RNA from lung tissue and transcription of *cramp*, *BD1*, *BD3*, *SP-A* and *SP-D*. At day 7 post influenza challenge, transcription of *cramp*, *bd1* and *sp-A* was significantly reduced in comparison to uninfected mice. In coinfecting mice, mRNA levels of *cramp* and *bd1* were strongly decreased. These findings are coherent with another study, which reported a downregulation of antimicrobial proteins and peptides in coinfecting versus single-infected lungs [304]. Interestingly, we found increased protein

expression of *cramp* in influenza and coinfecting mice compared to uninfected and single-infected ones. On one hand, increased cell death in the respiratory tract can lead to lower production levels of antimicrobial peptides and proteins. On the other hand, recruited neutrophils already contain *cramp* and release it, thus leading to a negative feedback, which inhibits *cramp* transcription and helps to prevent host cell damage [305].

It is well-known that influenza infections lead to increased oxidative stress in the LRT [306]. To study the effect of influenza-mediated ROS production by immune cells on pneumococcal virulence, we determined concentrations of  $H_2O_2$  in BAL of uninfected and influenza-challenged mice. After infection with influenza, we detected higher  $H_2O_2$  levels in BAL than in uninfected mice. Since pneumococcal resistance to ROS is regulated by the serine protease HtrA [234], we measured *htrA* transcription and found it to be increased in response to influenza-infected BAL compared to uninfected BAL. Hence we suggest that influenza leads to enhanced stress in form of denatured proteins and increased ROS in the LRT, which upregulates pneumococcal HtrA. Furthermore, in paper I we found a Sia-mediated upregulation of HtrA. Influenza infections increase the level of free Sia in the airway of infected mice [171], and thus could also mediate HtrA upregulation in pneumococci incubated in BAL of influenza-infected animals.

Genes involved in the regulation of oxidative stress have been shown to be important during coinfections of influenza and *H. influenzae* [307]. However, we did not find any study, which examined the role of HtrA in influenza/pneumococcal coinfections. The serine protease HtrA is surface exposed [241] and a well-known pneumococcal virulence factor [248]. It is demonstrated to have chaperone and protease functions, as it aids protein folding at lower temperatures, and degrades misfolded proteins under stress conditions [242]. Moreover, it has been demonstrated to degrade E-cadherin, thus fostering bacterial invasion across epithelial barriers [246, 247]. We detected higher protein concentrations and increased numbers of dead cells in the LRT of influenza infected mice at day 7 post infections. Thus, we hypothesized that pneumococci might metabolize and grow on denatured proteins and cell debris. To test the role of HtrA in a coinfection setting, we infected mock and influenza-challenged mice with TIGR4 $\Delta$ *htrA* and observed severely reduced bacterial numbers in the LRT of coinfecting in contrast to single-infected mice.

The TCS CiaRH is reported to control transcription of *htrA* *in vitro* [218, 219]. In contrast to infections with TIGR4 $\Delta$ *htrA*, infections with a *ciaR*-deficient mutant led to similar bacterial numbers as challenges with wt TIGR4. Moreover, our results suggest that HtrA is required to cope with stress in the LRT post influenza challenge and that *htrA* transcription can be controlled independently of CiaR in the lungs of coinfecting mice. Interestingly, RR09 has been implicated in pneumococcal virulence, as well as in regulation of HtrA expression [308]. Future studies will show, if RR09 is responsible for the upregulation of HtrA *in vivo* in the absence of CiaR.

In summary, our data suggest that influenza-mediated changes of the host environment in the LRT support secondary pneumococcal infections by promoting pneumococcal adaptation and

upregulating pneumococcal virulence factors, like HtrA. Moreover, HtrA facilitates pneumococcal outgrowth during influenza/pneumococcal coinfections, which indicates its importance as a therapeutic target.

## 4.4 PAPER IV

### **Toll-like receptor 3/TRIF-dependent IL-12p70 secretion mediated by *Streptococcus pneumoniae* RNA and its priming by influenza A virus coinfection in human dendritic cells**

The last study of this thesis describes the recognition of pneumococcal RNA by TLR3 and presents a mechanism for the increased IL-12 production in DCs upon influenza/pneumococcal coinfection.

DCs are shown to be the main producers of IL-12p70. After pneumococcal infections, IL-12 activates T<sub>H</sub>1-cells and stimulates the production of IFN- $\gamma$  by other immune cells [115-117]. IL-12 plays an important role *in vivo*, as its exogenous administration improved the disease outcome in a murine model of pneumococcal pneumonia [309]. Moreover, IL-12 deficiency in humans was associated with recurrent pneumococcal infections [310]. To examine the role of this cytokine in pneumococcal infections in monocyte-derived DCs, we challenged these cells with a low infection dose of T4R. A MOI of 1 did not affect cytotoxicity but evoked a secretion of IL-12p70 by DCs, which was dependent on pneumococcal uptake.

Next, we aimed at studying the role of TRIF in pneumococcal infections of DCs. TLR3 and TLR4 are the only PRRs that signal via TRIF, while other TLRs use MyD88 as adapter molecule to stimulate host responses [61, 62]. We silenced TRIF in DCs using small interfering RNAs and challenged the cells with T4R to measure IL-12p70 production. RNA as well as protein levels of IL-12 were decreased. Hence, DCs sense pneumococci via TRIF-mediated TLR-signalling.

To determine the role of TLR4 in pneumococcal sensing by DCs, we next silenced TLR4 in these cells and measured IL-12p70 secretion. While there are studies implicating pneumococcal toxin pneumolysin in the activation of TLR4 [67, 311-313], other research groups have observed a TLR4-independent sensing of pneumolysin [71, 251, 314]. Our data indicate that pneumococci can activate DCs independently of TLR4 expression in this model, as IL-12 transcription and secretion were not affected by inactivation of this PRR.

Next, we silenced TLR3 in DCs and measured IL-12p70 production after pneumococcal challenge. Interestingly, we found a decrease of this cytokine on transcriptional as well as on protein level. To confirm this result, we made use of a TLR3/dsRNA complex inhibitor, which reduced the secretion of IL-12. The capsule did not interfere with this result, as we observed a downregulation of IL-12 release even after challenge with wt TIGR4. These results show that DCs recognize pneumococci via the endosomal receptor TLR3.

A previous study by our group reported an IAV-mediated type I IFN production, which led to an enhanced secretion of IL-12 by neighbouring DCs [315]. Moreover, viral infections as well as type I IFN are known to upregulate TLR3 expression [316, 317]. To explore whether IAV upregulates TLR3 and consequently leads to an increase of IL-12p70 secretion in



influenza/pneumococcal coinfections, we measured mRNA levels of TLR3 in DCs challenged with influenza alone and after influenza/pneumococcal coinfection. Indeed, TLR3 expression was enhanced in IAV infected and coinfecting DCs. Moreover, we tested IL-12p40 transcription and found it to be upregulated in T4R single-infected and even more in IAV/T4R coinfecting DCs. However, treatment with IFN- $\alpha$  alone only had a minor effect on TLR3 expression, suggesting that additional factors other than IFN- $\alpha$  might contribute to the influenza-induced TLR3 upregulation.

TLR3 is known to recognize viral dsRNA [62]. However, there are recent reports describing TLR3-mediated sensing of bacterial RNA [318-320]. To test whether pneumococcal RNA is sufficient to induce TLR3-mediated IL-12p70 production, we isolated RNA from T4R and transfected it into DCs. We observed an induction of IL-12p70 secretion upon transfection with total RNA, which was dependent on TLR3 and completely abolished after RNase-treatment. Furthermore, we confirmed RNA-mediated TLR3 activation in HEK293 cells expressing TLR3.

Additionally, we killed pneumococci by heat exposure and UV radiation and found the latter condition to induce IL-12p70 secretion. We determined the RNA quality after the two treatments and found the RNA to be degraded after heat treatment, while UV radiation maintained RNA integrity. Moreover, our data show that RNase treatment of UV-killed pneumococci reduced cytokine secretion in a dose-dependent manner, while DNase treatment and LPS transfection in combination with RNase did not affect IL-12 production.

In summary our data indicate that pneumococci are taken up into DCs and degraded inside the endosome. Released pneumococcal RNA can subsequently be sensed by TLR3 and induces secretion of IL12-p70. Furthermore, our results give a possible explanation for the increased cytokine secretion in influenza/pneumococcal coinfection. The data suggest that infections with influenza lead to the secretion of type I IFN and other factors, which enhances TLR3 expression by DCs. Consequently, the higher level of TLR3 mediate increased IL12-p70 production in response to secondary pneumococcal infections.

## 5 CONCLUDING REMARKS

*S. pneumoniae* is a major threat to human health, especially of very young, elderly and immunocompromised individuals. Even though vaccines and antibiotics are available today, the emergence of non-vaccine types in IPD and antibiotic resistance in pneumococci demand for the development of improved vaccines and therapeutics. Consequently, a comprehensive understanding of pneumococcal-host interactions is required.

Immediately after entering the host, pulmonary pathogens, like *S. pneumoniae* or IAV, come in contact with glycans, which serve as receptors for adhesion [166] and thus present a prerequisite for pathogenesis. In the case of pneumococci, sugars of the glycan strand can also act as carbon source [172, 197, 293]. *S. pneumoniae* allocates more than 30% of all transporters to the uptake of carbohydrates [188], which points towards the importance of carbohydrates in pneumococcal colonization and disease. Sialic acid, the terminal sugar of the glycan strand, has been implicated in pneumococcal metabolism but has also been demonstrated to serve as a signal for pneumococcal disease [170, 172]. However, there is also another aspect of glycan-based host-pathogen interactions, as most PAMPs are demonstrated to be microbial glycan structures, and thus activate innate immune responses [181].

In paper I and II, we study sialic acid-mediated aspects of pneumococcal host tropism. In paper I, we demonstrate that pneumococci upregulate several virulence factors in response to Neu5Ac, the sialic acid mainly abundant in humans, compared with Neu5Gc, which cannot be synthesized by humans. Moreover, we show that sialic acid or its metabolic products are sensed involving the pneumococcal response regulator CiaR. In paper II, we show an increased pneumococcal adhesion rate to Neu5Ac– compared with Neu5Gc-containing host glycans. Moreover, we present data indicating a role of pneumococcal cytotoxins in increased immune evasion by pneumococci in response to Neu5Ac compared with Neu5Gc.

In paper III and IV, we study pneumococcal-host interactions after prior influenza infections, since they account for severe disease outcomes. In paper III, we determine influenza-induced changes in the LRT of mice and investigate pneumococcal adaptations to this environment. In paper IV, we show that pneumococcal RNA can be sensed by TLR3 in DCs. Moreover, in a coinfection setting, influenza infections increase TLR3 expression with the help of type I interferon. Thus upon secondary pneumococcal challenge, enhanced TLR3 expression upregulates IL-12 production.

In summary, we show that glycans play a pivotal role in pneumococcal pathogenesis. The studies included in this work give insights into alterations of pneumococcal-host interactions caused by differences in the sialic acid profile of humans in comparison to other mammals, as well as by preceding viral infections. Moreover, this thesis describes the activation of the innate immune system by pneumococcal glycan-based structures, like RNA. These findings will hopefully add to the improvement of current treatment options and to the development of new vaccines.

## 6 ACKNOWLEDGEMENTS

There are many people who supported me during my time as a PhD student and who made it truly unforgettable. I would especially like to thank...

...my main supervisor **Birgitta Henriques-Normark**. I feel privileged that I got the chance to work on this exciting project. Thank you for your enormous support, for encouraging me to pursue my own scientific hypotheses but also for your help, when projects did not work out as expected. I am also really grateful that I was allowed to attend all those interesting courses and conferences.

...my co-supervisor **Staffan Normark**. Thank you for allocating time for long meetings in phases, in which the project seemed to be stuck, for seeing the bigger picture of results and for your scientific input, which often added a new and exciting twist.

...my co-supervisor **Jonas Löfling**. Thank you for letting me join this fantastic project and for introducing me to the “sugar world”. I really appreciate that you always offered your help if needed, even though you left the group quite a while ago.

...**Ajit Varki** and **Victor Nizet** from the University of California, San Diego. I am really grateful for the weeks I spent in your research groups to learn new techniques. Everyone was really welcoming and helpful. I would like to thank you in particular for the valuable input you gave on the project and manuscript.

...and all **co-authors** on the publications and manuscripts for letting me join their interesting projects or for supporting my project with experiments and/or constructive feedback.

...**Laura**, for giving me the chance to join your exciting work, which became part of this thesis. Thank you for being calm when others are not, for nice chats over coffee, for the great time we had on several summer schools and conferences, and for sharing the enthusiasm for running. I am looking forward to the half marathon 2018.

...**Vicky**, for introducing me to the exciting field of influenza/pneumococcal coinfection, for nice chats at work and also outside work, for your help during long days in the animal house, for the great memories we share of our trips to India or the US, for your honesty and for ensuring that I do not forget science during the last months of leave.

...**Anuj**, for being a great person and pleasant office mate (not only for providing a huge selection of afternoon snacks), for teaching me how to cook delicious Indian food, for the help during our many movings, and for the good times we had during Christmas in Germany.

... **Elisabeth**, for being incredibly kind. I really enjoyed being your office neighbour. **Mario**, **Hannes** and **Jens**, for creating a fun atmosphere in the office and for keeping me up to date on what’s trending on the internet among young people.

...all past and present members of the **BHN**, as well as the **Rhen** and **Loh** group, for fun coffee breaks, interesting discussions in lab meetings or during lunch breaks and for creating a great work environment.

...**Ilias**, for being a great travel companion, for relaxing lunches at Nanna and for being sincere and always helpful. **Martin**, for sharing your “pipettpojke” with me and your patience in teaching me *in vivo* techniques. **Murat** and **Alice** - You left the group for new adventures but I am always happy to see or read what you are up to.

...**Anna-Karin** and **Torunn** and all other staff members of the animal facility, for being indispensable during *in vivo* studies. Thank you for all your help and efforts!

...all organizers and teaching assistants of the **DSM2.1** Laboratory course in Medical Microbiology, in particular **Dae-Ho**, **Patricia** and **Sara**. Thank you for the great experience. I learned so much and really enjoyed being part of the team!

...and of course all the important people outside my work life:

...**Vilma**, I still remember the first time meeting you during our master theses at MTC. I felt an immediate connection and it did not change ever since. Thank you for being a good listener, a problem solver and an incredible friend who I can always trust.

...**Cécile**, for your directness and honesty; **Ale**, for the great time we had on concerts in Debaser; **Andrea & Steffi**, for fun nights out and inviting me to your amazing wedding party in Italy; **Haythem**, for being a really kind person; **Mustafa**, for fun discussions about all kinds of topics, **Susi K**, für deine angenehme Art und deinen norddeutschen Humor; und **Tina**, für die tolle Zeit, die wir im Jägargatan hatten und für großartige Abende im Nachtleben von Stockholm.

...**Anni**, **Celli**, **Bine**, **Chrissy**, **Jessi**, **Marius** und **Annie** – Ihr seid ein wichtiger Teil meines Lebens und einer der Gründe, warum ich immer wieder gern in die Heimat fahre. Danke, dass es sich immer so anfühlt, als ob ich nie weg war, wenn wir uns sehen. **Susi** und **Björn**, **Krischi** und **Manu**, **Susi** und **Henner**. Danke, für die tollen Skype-Abende, eure Besuche hier in Stockholm oder die schöne Zeit, die wir haben, wenn wir in Magdeburg sind. Ich freue mich schon auf die aufregenden Jahre, die vor uns liegen. **Gino**, dafür, dass du ein toller Mitbewohner warst und für deine endlose Hilfsbereitschaft, nicht nur in IT-Fragen.

...**Christians Familie**, für eure Warmherzigkeit, Hilfsbereitschaft und Unterstützung.

...natürlich meiner Familie. **Mutti**, **Vati**, **Felix**, **Guido**, **Nici**, **Fynni**, **Elinchen**, **Oma** und **Opa** – Danke für eure unendliche Unterstützung, schon während des Studiums, und dafür, dass ihr immer an mich geglaubt habt.

...und **Christian**, dafür, dass du der beste Freund bist, den ich mir wünschen könnte. Danke, für den Halt, den du mir gegeben hast, nicht nur während der letzten Monate.

## 7 REFERENCES

1. Austrian, R., *The enduring pneumococcus: unfinished business and opportunities for the future*. Microb Drug Resist, 1997. **3**(2): p. 111-5.
2. Pasteur, L., L. Chamberland, and E. Roux, *Sur une maladie nouvelle provoquée par la salive d'un enfant mort la rage*. C.R Acad XCLI, 1881. **92**: p. 159-165.
3. Watson, D.A., et al., *A brief history of the pneumococcus in biomedical research: a panoply of scientific discovery*. Clin Infect Dis, 1993. **17**(5): p. 913-24.
4. Geno, K.A., et al., *Pneumococcal Capsules and Their Types: Past, Present, and Future*. Clin Microbiol Rev, 2015. **28**(3): p. 871-99.
5. Martin-Galiano, A.J., et al., *Genetic characterization of optochin-susceptible viridans group streptococci*. Antimicrob Agents Chemother, 2003. **47**(10): p. 3187-94.
6. Moore, H.F., *The Action of Ethylhydrocuprein (Optochin) on Type Strains of Pneumococci in Vitro and in Vivo, and on Some Other Microorganisms in Vitro*. J Exp Med, 1915. **22**(3): p. 269-85.
7. Aguiar, S.I., et al., *Emergence of optochin resistance among Streptococcus pneumoniae in Portugal*. Microb Drug Resist, 2006. **12**(4): p. 239-45.
8. Griffith, F., *The Significance of Pneumococcal Types*. J Hyg (Lond), 1928. **27**(2): p. 113-59.
9. Avery, O.T., C.M. Macleod, and M. McCarty, *Studies on the Chemical Nature of the Substance Inducing Transformation of Pneumococcal Types : Induction of Transformation by a Desoxyribonucleic Acid Fraction Isolated from Pneumococcus Type Iii*. J Exp Med, 1944. **79**(2): p. 137-58.
10. Henriques-Normark, B. and E.I. Tuomanen, *The pneumococcus: epidemiology, microbiology, and pathogenesis*. Cold Spring Harb Perspect Med, 2013. **3**(7).
11. O'Brien, K.L., et al., *Burden of disease caused by Streptococcus pneumoniae in children younger than 5 years: global estimates*. Lancet, 2009. **374**(9693): p. 893-902.
12. Kadioglu, A., et al., *The role of Streptococcus pneumoniae virulence factors in host respiratory colonization and disease*. Nature reviews. Microbiology, 2008. **6**(4): p. 288-301.
13. Mato, R., et al., *Natural history of drug-resistant clones of Streptococcus pneumoniae colonizing healthy children in Portugal*. Microbial drug resistance, 2005. **11**(4): p. 309-22.
14. Rey, L.C., et al., *Antimicrobial susceptibility and serotypes of nasopharyngeal Streptococcus pneumoniae in children with pneumonia and in children attending day-care centres in Fortaleza, Brazil*. International journal of antimicrobial agents, 2002. **20**(2): p. 86-92.
15. Nunes, S., et al., *Trends in drug resistance, serotypes, and molecular types of Streptococcus pneumoniae colonizing preschool-age children attending day care centers in Lisbon, Portugal: a summary of 4 years of annual surveillance*. J Clin Microbiol, 2005. **43**(3): p. 1285-93.

16. Wyllie, A.L., et al., *Molecular surveillance on Streptococcus pneumoniae carriage in non-elderly adults; little evidence for pneumococcal circulation independent from the reservoir in children*. Sci Rep, 2016. **6**: p. 34888.
17. Rovers, M.M., et al., *Otitis media*. Lancet, 2004. **363**(9407): p. 465-73.
18. Celind, J., L. Sodermark, and O. Hjalmarson, *Adherence to treatment guidelines for acute otitis media in children. The necessity of an effective strategy of guideline implementation*. Int J Pediatr Otorhinolaryngol, 2014. **78**(7): p. 1128-32.
19. Bluestone, C.D., *Clinical course, complications and sequelae of acute otitis media*. Pediatr Infect Dis J, 2000. **19**(5 Suppl): p. S37-46.
20. Brook, I., *Microbiology and choice of antimicrobial therapy for acute sinusitis complicated by subperiosteal abscess in children*. Int J Pediatr Otorhinolaryngol, 2016. **84**: p. 21-6.
21. Rosenfeld, R.M., et al., *Clinical practice guideline (update): Adult Sinusitis Executive Summary*. Otolaryngol Head Neck Surg, 2015. **152**(4): p. 598-609.
22. Rose, M.A., et al., *The burden of invasive pneumococcal disease in children with underlying risk factors in North America and Europe*. Int J Clin Pract, 2014. **68**(1): p. 8-19.
23. Torres, A., et al., *The aetiology and antibiotic management of community-acquired pneumonia in adults in Europe: a literature review*. Eur J Clin Microbiol Infect Dis, 2014. **33**(7): p. 1065-79.
24. Feldman, C. and R. Anderson, *Epidemiology, virulence factors and management of the pneumococcus*. F1000Res, 2016. **5**: p. 2320.
25. Rudan, I., et al., *Epidemiology and etiology of childhood pneumonia*. Bull World Health Organ, 2008. **86**(5): p. 408-16.
26. Maraqa, N.F., *Pneumococcal infections*. Pediatr Rev, 2014. **35**(7): p. 299-310.
27. Angus, D.C., et al., *Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care*. Crit Care Med, 2001. **29**(7): p. 1303-10.
28. Schuchat, A., et al., *Bacterial meningitis in the United States in 1995. Active Surveillance Team*. N Engl J Med, 1997. **337**(14): p. 970-6.
29. Rajasingham, C.R., et al., *Serious neurologic sequelae in cases of meningitis arising from infection by conjugate vaccine-related and nonvaccine-related serogroups of Streptococcus pneumoniae*. Pediatr Infect Dis J, 2008. **27**(9): p. 771-5.
30. Hoogman, M., et al., *Cognitive outcome in adults after bacterial meningitis*. J Neurol Neurosurg Psychiatry, 2007. **78**(10): p. 1092-6.
31. Weisfelt, M., et al., *Clinical features, complications, and outcome in adults with pneumococcal meningitis: a prospective case series*. Lancet Neurol, 2006. **5**(2): p. 123-9.
32. Robinson, K.A., et al., *Epidemiology of invasive Streptococcus pneumoniae infections in the United States, 1995-1998: Opportunities for prevention in the conjugate vaccine era*. JAMA, 2001. **285**(13): p. 1729-35.

33. Bogaert, D., R. De Groot, and P.W. Hermans, *Streptococcus pneumoniae* colonisation: the key to pneumococcal disease. *Lancet Infect Dis*, 2004. **4**(3): p. 144-54.
34. Falsey, A.R., et al., *Bacterial complications of respiratory tract viral illness: a comprehensive evaluation*. *J Infect Dis*, 2013. **208**(3): p. 432-41.
35. McCullers, J.A., *The co-pathogenesis of influenza viruses with bacteria in the lung*. *Nat Rev Microbiol*, 2014. **12**(4): p. 252-62.
36. Morens, D.M., J.K. Taubenberger, and A.S. Fauci, *Predominant role of bacterial pneumonia as a cause of death in pandemic influenza: implications for pandemic influenza preparedness*. *J Infect Dis*, 2008. **198**(7): p. 962-70.
37. DiMenna, L.J. and H.C. Ertl, *Pandemic influenza vaccines*. *Curr Top Microbiol Immunol*, 2009. **333**: p. 291-321.
38. Schwarzmann, S.W., et al., *Bacterial pneumonia during the Hong Kong influenza epidemic of 1968-1969*. *Arch Intern Med*, 1971. **127**(6): p. 1037-41.
39. Martin, C.M., et al., *Asian influenza A in Boston, 1957-1958. I. Observations in thirty-two influenza-associated fatal cases*. *AMA Arch Intern Med*, 1959. **103**(4): p. 515-31.
40. Rice, T.W., et al., *Critical illness from 2009 pandemic influenza A virus and bacterial coinfection in the United States*. *Crit Care Med*, 2012. **40**(5): p. 1487-98.
41. Hakenbeck, R., et al., *Molecular mechanisms of beta-lactam resistance in Streptococcus pneumoniae*. *Future Microbiol*, 2012. **7**(3): p. 395-410.
42. Kislak, J.W., et al., *Susceptibility of pneumococci to nine antibiotics*. *Am J Med Sci*, 1965. **250**(3): p. 261-8.
43. *European Centre for Disease Prevention and Control. Antimicrobial resistance surveillance in Europe 2013. Annual Report of the European Antimicrobial Resistance Surveillance Network (EARS-Net)*. 2014: Stockholm.
44. Schroeder, M.R. and D.S. Stephens, *Macrolide Resistance in Streptococcus pneumoniae*. *Front Cell Infect Microbiol*, 2016. **6**: p. 98.
45. Felmingham, D., R. Canton, and S.G. Jenkins, *Regional trends in beta-lactam, macrolide, fluoroquinolone and telithromycin resistance among Streptococcus pneumoniae isolates 2001-2004*. *J Infect*, 2007. **55**(2): p. 111-8.
46. *Antibiotic resistance threats in the United States, 2013*, CDC, Editor. 2013: USA.
47. Feldman, C. and R. Anderson, *Review: current and new generation pneumococcal vaccines*. *J Infect*, 2014. **69**(4): p. 309-25.
48. Koskela, M., et al., *First and second dose antibody responses to pneumococcal polysaccharide vaccine in infants*. *Pediatr Infect Dis*, 1986. **5**(1): p. 45-50.
49. Reinert, R.R., P. Paradiso, and B. Fritzell, *Advances in pneumococcal vaccines: the 13-valent pneumococcal conjugate vaccine received market authorization in Europe*. *Expert Rev Vaccines*, 2010. **9**(3): p. 229-36.
50. Paradiso, P.R., *Pneumococcal conjugate vaccine for adults: a new paradigm*. *Clin Infect Dis*, 2012. **55**(2): p. 259-64.

51. Rose, M. and S. Zielen, *Impact of infant immunization programs with pneumococcal conjugate vaccine in Europe*. Expert Rev Vaccines, 2009. **8**(10): p. 1351-64.
52. *Vaccinationsstatistik från barnavårdscentralerna 2016, gällande barn födda 2013*, Folkhälsomyndigheten, Editor. 2016: Stockholm.
53. Galanis, I., et al., *Effects of PCV7 and PCV13 on invasive pneumococcal disease and carriage in Stockholm, Sweden*. Eur Respir J, 2016. **47**(4): p. 1208-18.
54. Browall, S., et al., *Clinical manifestations of invasive pneumococcal disease by vaccine and non-vaccine types*. Eur Respir J, 2014. **44**(6): p. 1646-57.
55. Feikin, D.R., et al., *Serotype-specific changes in invasive pneumococcal disease after pneumococcal conjugate vaccine introduction: a pooled analysis of multiple surveillance sites*. PLoS Med, 2013. **10**(9): p. e1001517.
56. Briles, D.E., et al., *The potential for using protein vaccines to protect against otitis media caused by Streptococcus pneumoniae*. Vaccine, 2000. **19 Suppl 1**: p. S87-95.
57. Ogunniyi, A.D., et al., *Immunization of mice with combinations of pneumococcal virulence proteins elicits enhanced protection against challenge with Streptococcus pneumoniae*. Infect Immun, 2000. **68**(5): p. 3028-33.
58. Ogunniyi, A.D., P. Giammarinaro, and J.C. Paton, *The genes encoding virulence-associated proteins and the capsule of Streptococcus pneumoniae are upregulated and differentially expressed in vivo*. Microbiology, 2002. **148**(Pt 7): p. 2045-53.
59. Hernani Mde, L., et al., *Nasal immunization of mice with Lactobacillus casei expressing the pneumococcal surface protein C primes the immune system and decreases pneumococcal nasopharyngeal colonization in mice*. FEMS Immunol Med Microbiol, 2011. **62**(3): p. 263-72.
60. Medzhitov, R. and C. Janeway, Jr., *Innate immunity*. N Engl J Med, 2000. **343**(5): p. 338-44.
61. Kawai, T. and S. Akira, *The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors*. Nat Immunol, 2010. **11**(5): p. 373-84.
62. Akira, S., S. Uematsu, and O. Takeuchi, *Pathogen recognition and innate immunity*. Cell, 2006. **124**(4): p. 783-801.
63. Chen, G., et al., *NOD-like receptors: role in innate immunity and inflammatory disease*. Annu Rev Pathol, 2009. **4**: p. 365-98.
64. Mariathasan, S., et al., *Cryopyrin activates the inflammasome in response to toxins and ATP*. Nature, 2006. **440**(7081): p. 228-32.
65. Yoshimura, A., et al., *Cutting edge: recognition of Gram-positive bacterial cell wall components by the innate immune system occurs via Toll-like receptor 2*. J Immunol, 1999. **163**(1): p. 1-5.
66. Albiger, B., et al., *Toll-like receptor 9 acts at an early stage in host defence against pneumococcal infection*. Cell Microbiol, 2007. **9**(3): p. 633-44.
67. Malley, R., et al., *Recognition of pneumolysin by Toll-like receptor 4 confers resistance to pneumococcal infection*. Proc Natl Acad Sci U S A, 2003. **100**(4): p. 1966-71.



68. Lemon, J.K. and J.N. Weiser, *Degradation products of the extracellular pathogen Streptococcus pneumoniae access the cytosol via its pore-forming toxin*. MBio, 2015. **6**(1).
69. Davis, K.M., S. Nakamura, and J.N. Weiser, *Nod2 sensing of lysozyme-digested peptidoglycan promotes macrophage recruitment and clearance of S. pneumoniae colonization in mice*. J Clin Invest, 2011. **121**(9): p. 3666-76.
70. Witzenrath, M., et al., *The NLRP3 inflammasome is differentially activated by pneumolysin variants and contributes to host defense in pneumococcal pneumonia*. J Immunol, 2011. **187**(1): p. 434-40.
71. McNeela, E.A., et al., *Pneumolysin activates the NLRP3 inflammasome and promotes proinflammatory cytokines independently of TLR4*. PLoS Pathog, 2010. **6**(11): p. e1001191.
72. Koppe, U., N. Suttorp, and B. Opitz, *Recognition of Streptococcus pneumoniae by the innate immune system*. Cellular microbiology, 2012. **14**(4): p. 460-6.
73. Sprangers, S., T.J. de Vries, and V. Everts, *Monocyte Heterogeneity: Consequences for Monocyte-Derived Immune Cells*. J Immunol Res, 2016. **2016**: p. 1475435.
74. Monick, M.M. and G.W. Hunninghake, *Activation of second messenger pathways in alveolar macrophages by endotoxin*. Eur Respir J, 2002. **20**(1): p. 210-22.
75. Martin, T.R. and C.W. Frevert, *Innate immunity in the lungs*. Proc Am Thorac Soc, 2005. **2**(5): p. 403-11.
76. Thomas, C.A., et al., *Protection from lethal gram-positive infection by macrophage scavenger receptor-dependent phagocytosis*. J Exp Med, 2000. **191**(1): p. 147-56.
77. Greenberg, S. and S. Grinstein, *Phagocytosis and innate immunity*. Curr Opin Immunol, 2002. **14**(1): p. 136-45.
78. Kraal, G., et al., *The macrophage receptor MARCO*. Microbes Infect, 2000. **2**(3): p. 313-6.
79. Gwinn, M.R. and V. Vallyathan, *Respiratory burst: role in signal transduction in alveolar macrophages*. J Toxicol Environ Health B Crit Rev, 2006. **9**(1): p. 27-39.
80. Arredouani, M., et al., *The scavenger receptor MARCO is required for lung defense against pneumococcal pneumonia and inhaled particles*. J Exp Med, 2004. **200**(2): p. 267-72.
81. Kang, Y.S., et al., *The C-type lectin SIGN-R1 mediates uptake of the capsular polysaccharide of Streptococcus pneumoniae in the marginal zone of mouse spleen*. Proc Natl Acad Sci U S A, 2004. **101**(1): p. 215-20.
82. Orrskog, S., et al., *Pilus adhesin RrgA interacts with complement receptor 3, thereby affecting macrophage function and systemic pneumococcal disease*. mBio, 2012. **4**(1): p. e00535-12.
83. Nelson, A.L., et al., *Capsule enhances pneumococcal colonization by limiting mucus-mediated clearance*. Infect Immun, 2007. **75**(1): p. 83-90.
84. Marriott, H.M., et al., *Contrasting roles for reactive oxygen species and nitric oxide in the innate response to pulmonary infection with Streptococcus pneumoniae*. Vaccine, 2007. **25**(13): p. 2485-90.

85. Bainton, D.F., J.L. Ulliyot, and M.G. Farquhar, *The development of neutrophilic polymorphonuclear leukocytes in human bone marrow*. J Exp Med, 1971. **134**(4): p. 907-34.
86. McCullers, J.A. and E.I. Tuomanen, *Molecular pathogenesis of pneumococcal pneumonia*. Front Biosci, 2001. **6**: p. D877-89.
87. Jones, M.R., et al., *Lung NF-kappaB activation and neutrophil recruitment require IL-1 and TNF receptor signaling during pneumococcal pneumonia*. J Immunol, 2005. **175**(11): p. 7530-5.
88. Mayer-Scholl, A., P. Averhoff, and A. Zychlinsky, *How do neutrophils and pathogens interact?* Curr Opin Microbiol, 2004. **7**(1): p. 62-6.
89. Brinkmann, V., et al., *Neutrophil extracellular traps kill bacteria*. Science, 2004. **303**(5663): p. 1532-5.
90. Standish, A.J. and J.N. Weiser, *Human neutrophils kill Streptococcus pneumoniae via serine proteases*. J Immunol, 2009. **183**(4): p. 2602-9.
91. Marriott, H.M., et al., *Reactive oxygen species regulate neutrophil recruitment and survival in pneumococcal pneumonia*. Am J Respir Crit Care Med, 2008. **177**(8): p. 887-95.
92. Wartha, F., et al., *Neutrophil extracellular traps: casting the NET over pathogenesis*. Curr Opin Microbiol, 2007. **10**(1): p. 52-6.
93. Beiter, K., et al., *An endonuclease allows Streptococcus pneumoniae to escape from neutrophil extracellular traps*. Curr Biol, 2006. **16**(4): p. 401-7.
94. O'Keeffe, M., W.H. Mok, and K.J. Radford, *Human dendritic cell subsets and function in health and disease*. Cell Mol Life Sci, 2015. **72**(22): p. 4309-25.
95. Bajtaj, Z., et al., *Expression and role of Fc- and complement-receptors on human dendritic cells*. Immunol Lett, 2006. **104**(1-2): p. 46-52.
96. Banchereau, J., et al., *Immunobiology of dendritic cells*. Annu Rev Immunol, 2000. **18**: p. 767-811.
97. Brogden, K.A., *Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria?* Nat Rev Microbiol, 2005. **3**(3): p. 238-50.
98. Peschel, A. and H.G. Sahl, *The co-evolution of host cationic antimicrobial peptides and microbial resistance*. Nat Rev Microbiol, 2006. **4**(7): p. 529-36.
99. Hiemstra, P.S., et al., *Antimicrobial Peptides and Innate Lung Defenses: Role in Infectious and Noninfectious Lung Diseases and Therapeutic Applications*. Chest, 2016. **149**(2): p. 545-51.
100. Hiemstra, P.S., P.B. McCray, Jr., and R. Bals, *The innate immune function of airway epithelial cells in inflammatory lung disease*. Eur Respir J, 2015. **45**(4): p. 1150-62.
101. Oren, Z., et al., *Structure and organization of the human antimicrobial peptide LL-37 in phospholipid membranes: relevance to the molecular basis for its non-cell-selective activity*. Biochem J, 1999. **341** ( Pt 3): p. 501-13.
102. Gallo, R.L., et al., *Identification of CRAMP, a cathelin-related antimicrobial peptide expressed in the embryonic and adult mouse*. J Biol Chem, 1997. **272**(20): p. 13088-93.

103. Lehrer, R.I. and T. Ganz, *Defensins of vertebrate animals*. Curr Opin Immunol, 2002. **14**(1): p. 96-102.
104. Moser, C., et al., *beta-Defensin 1 contributes to pulmonary innate immunity in mice*. Infect Immun, 2002. **70**(6): p. 3068-72.
105. Cruse, G., et al., *Human lung mast cells mediate pneumococcal cell death in response to activation by pneumolysin*. J Immunol, 2010. **184**(12): p. 7108-15.
106. Beiter, K., et al., *The capsule sensitizes Streptococcus pneumoniae to alpha-defensins human neutrophil proteins 1 to 3*. Infect Immun, 2008. **76**(8): p. 3710-6.
107. Scharf, S., et al., *Streptococcus pneumoniae induces human beta-defensin-2 and -3 in human lung epithelium*. Exp Lung Res, 2012. **38**(2): p. 100-10.
108. LeVine, A.M., et al., *Surfactant protein D enhances clearance of influenza A virus from the lung in vivo*. J Immunol, 2001. **167**(10): p. 5868-73.
109. Haagsman, H.P. and R.V. Diemel, *Surfactant-associated proteins: functions and structural variation*. Comp Biochem Physiol A Mol Integr Physiol, 2001. **129**(1): p. 91-108.
110. Haagsman, H.P., *Interactions of surfactant protein A with pathogens*. Biochim Biophys Acta, 1998. **1408**(2-3): p. 264-77.
111. Jounblat, R., et al., *Pneumococcal behavior and host responses during bronchopneumonia are affected differently by the cytolytic and complement-activating activities of pneumolysin*. Infect Immun, 2003. **71**(4): p. 1813-9.
112. Jounblat, R., et al., *The role of surfactant protein D in the colonisation of the respiratory tract and onset of bacteraemia during pneumococcal pneumonia*. Respir Res, 2005. **6**: p. 126.
113. Kuronuma, K., et al., *Pulmonary surfactant protein A augments the phagocytosis of Streptococcus pneumoniae by alveolar macrophages through a casein kinase 2-dependent increase of cell surface localization of scavenger receptor A*. J Biol Chem, 2004. **279**(20): p. 21421-30.
114. Broere, F.A., S.G.; Sitkovsky, M.V.; Van Eden, W., *T cell subsets and T cell-mediated immunity*. Principles of Immunopharmacology, 2011. **3**.
115. Yamamoto, N., et al., *Essential role for the p40 subunit of interleukin-12 in neutrophil-mediated early host defense against pulmonary infection with Streptococcus pneumoniae: involvement of interferon-gamma*. Microbes Infect, 2004. **6**(14): p. 1241-9.
116. Kemp, K., et al., *Pneumococcal infections in humans are associated with increased apoptosis and trafficking of type 1 cytokine-producing T cells*. Infect Immun, 2002. **70**(9): p. 5019-25.
117. Rubins, J.B. and C. Pomeroy, *Role of gamma interferon in the pathogenesis of bacteremic pneumococcal pneumonia*. Infect Immun, 1997. **65**(7): p. 2975-7.
118. Hutter, J. and B. Lepenies, *Carbohydrate-Based Vaccines: An Overview*. Methods Mol Biol, 2015. **1331**: p. 1-10.
119. Parker, D.C., *The functions of antigen recognition in T cell-dependent B cell activation*. Semin Immunol, 1993. **5**(6): p. 413-20.

120. Rambourg, A. and C.P. Leblond, *Electron microscope observations on the carbohydrate-rich cell coat present at the surface of cells in the rat*. The Journal of cell biology, 1967. **32**(1): p. 27-53.
121. Varki, A., *Essentials of Glycobiology*. 1999, New York: Cold Spring Harbor Laboratory Press.
122. Nothhaft, H. and C.M. Szymanski, *Protein glycosylation in bacteria: sweeter than ever*. Nat Rev Microbiol, 2010. **8**(11): p. 765-78.
123. Varki, A., *Glycan-based interactions involving vertebrate sialic-acid-recognizing proteins*. Nature, 2007. **446**(7139): p. 1023-9.
124. Tra, V.N. and D.H. Dube, *Glycans in pathogenic bacteria--potential for targeted covalent therapeutics and imaging agents*. Chem Commun (Camb), 2014. **50**(36): p. 4659-73.
125. Blix, G., *Über die Kohlenhydratgruppen aus dem Submaxillarmucin*. Hoppe Seylers Z Physiol Chem, 1936. **240**(1-2): p. 43-54.
126. Klenk, E., *Neuraminsäure, das Spaltprodukt eines neuen Gehirnlipoids*. Hoppe Seylers Z Physiol Chem, 1941. **268**(1-2): p. 50-58.
127. Blix, F.G., A. Gottschalk, and E. Klenk, *Proposed nomenclature in the field of neuraminic and sialic acids*. Nature, 1957. **179**(4569): p. 1088.
128. Schauer, R., *Sialic acids as regulators of molecular and cellular interactions*. Current opinion in structural biology, 2009. **19**(5): p. 507-14.
129. Vimr, E. and C. Lichtensteiger, *To sialylate, or not to sialylate: that is the question*. Trends in microbiology, 2002. **10**(6): p. 254-7.
130. Malykh, Y.N., et al., *The presence of N-acetylneuraminic acid in Malpighian tubules of larvae of the cicada Philaenus spumarius*. Glycoconj J, 1999. **16**(11): p. 731-9.
131. Roth, J., et al., *Occurrence of sialic acids in Drosophila melanogaster*. Science, 1992. **256**(5057): p. 673-5.
132. Shaw, L. and R. Schauer, *Detection of CMP-N-acetylneuraminic acid hydroxylase activity in fractionated mouse liver*. The Biochemical journal, 1989. **263**(2): p. 355-63.
133. Chou, H.H., et al., *A mutation in human CMP-sialic acid hydroxylase occurred after the Homo-Pan divergence*. Proceedings of the National Academy of Sciences of the United States of America, 1998. **95**(20): p. 11751-6.
134. Muchmore, E.A., S. Diaz, and A. Varki, *A structural difference between the cell surfaces of humans and the great apes*. American journal of physical anthropology, 1998. **107**(2): p. 187-98.
135. Okerblom, J.J., et al., *Loss of CMAH during Human Evolution Primed the Monocyte-Macrophage Lineage toward a More Inflammatory and Phagocytic State*. J Immunol, 2017. **198**(6): p. 2366-2373.
136. Ghaderi, D., et al., *Sexual selection by female immunity against paternal antigens can fix loss of function alleles*. Proc Natl Acad Sci U S A, 2011. **108**(43): p. 17743-8.

137. Varki, A. and P. Gagneux, *Human-specific evolution of sialic acid targets: explaining the malignant malaria mystery?* Proceedings of the National Academy of Sciences of the United States of America, 2009. **106**(35): p. 14739-40.
138. Springer, S.A., S.L. Diaz, and P. Gagneux, *Parallel evolution of a self-signal: humans and new world monkeys independently lost the cell surface sugar Neu5Gc.* Immunogenetics, 2014. **66**(11): p. 671-4.
139. Ng, P.S., et al., *Ferrets exclusively synthesize Neu5Ac and express naturally humanized influenza A virus receptors.* Nature communications, 2014. **5**: p. 5750.
140. Martin, M.J., et al., *Evolution of human-chimpanzee differences in malaria susceptibility: relationship to human genetic loss of N-glycolylneuraminic acid.* Proceedings of the National Academy of Sciences of the United States of America, 2005. **102**(36): p. 12819-24.
141. Ward, J.M. and E.J. Vallender, *The resurgence and genetic implications of New World primates in biomedical research.* Trends in genetics : TIG, 2012. **28**(12): p. 586-91.
142. Deng, L., et al., *Host adaptation of a bacterial toxin from the human pathogen Salmonella Typhi.* Cell, 2014. **159**(6): p. 1290-9.
143. Naito, Y., et al., *Germinal center marker GL7 probes activation-dependent repression of N-glycolylneuraminic acid, a sialic acid species involved in the negative modulation of B-cell activation.* Mol Cell Biol, 2007. **27**(8): p. 3008-22.
144. Buchlis, G., et al., *Enhanced T cell function in a mouse model of human glycosylation.* J Immunol, 2013. **191**(1): p. 228-37.
145. Soto, P.C., et al., *Relative over-reactivity of human versus chimpanzee lymphocytes: implications for the human diseases associated with immune activation.* J Immunol, 2010. **184**(8): p. 4185-95.
146. Tangvoranuntakul, P., et al., *Human uptake and incorporation of an immunogenic nonhuman dietary sialic acid.* Proceedings of the National Academy of Sciences of the United States of America, 2003. **100**(21): p. 12045-50.
147. Hedlund, M., et al., *Evidence for a human-specific mechanism for diet and antibody-mediated inflammation in carcinoma progression.* Proc Natl Acad Sci U S A, 2008. **105**(48): p. 18936-41.
148. Samraj, A.N., et al., *A red meat-derived glycan promotes inflammation and cancer progression.* Proc Natl Acad Sci U S A, 2015. **112**(2): p. 542-7.
149. Koch, B.E., J. Stougaard, and H.P. Spaink, *Keeping track of the growing number of biological functions of chitin and its interaction partners in biomedical research.* Glycobiology, 2015. **25**(5): p. 469-82.
150. McFarlane, H.E., A. Doring, and S. Persson, *The cell biology of cellulose synthesis.* Annu Rev Plant Biol, 2014. **65**: p. 69-94.
151. Gibson, R., et al., *Synthesis and infectivity of vesicular stomatitis virus containing nonglycosylated G protein.* Cell, 1978. **13**(4): p. 671-9.
152. Lillehoj, E.P., et al., *Cellular and molecular biology of airway mucins.* Int Rev Cell Mol Biol, 2013. **303**: p. 139-202.

153. Schwarzkopf, M., et al., *Sialylation is essential for early development in mice*. Proc Natl Acad Sci U S A, 2002. **99**(8): p. 5267-70.
154. Varki, A., *Sialic acids in human health and disease*. Trends Mol Med, 2008. **14**(8): p. 351-60.
155. McEver, R.P., *Selectins: initiators of leucocyte adhesion and signalling at the vascular wall*. Cardiovasc Res, 2015. **107**(3): p. 331-9.
156. Varki, A., *Since there are PAMPs and DAMPs, there must be SAMPs? Glycan "self-associated molecular patterns" dampen innate immunity, but pathogens can mimic them*. Glycobiology, 2011. **21**(9): p. 1121-4.
157. Crocker, P.R., J.C. Paulson, and A. Varki, *Siglecs and their roles in the immune system*. Nat Rev Immunol, 2007. **7**(4): p. 255-66.
158. Ashwell, G. and T. Kawasaki, *A protein from mammalian liver that specifically binds galactose-terminated glycoproteins*. Methods Enzymol, 1978. **50**: p. 287-8.
159. Hoffmeister, K.M. and H. Falet, *Platelet clearance by the hepatic Ashwell-Morrell receptor: mechanisms and biological significance*. Thromb Res, 2016. **141 Suppl 2**: p. S68-72.
160. Lillehoj, E.R. and K.C. Kim, *Airway mucus: its components and function*. Arch Pharm Res, 2002. **25**(6): p. 770-80.
161. Gendler, S.J. and A.P. Spicer, *Epithelial mucin genes*. Annu Rev Physiol, 1995. **57**: p. 607-34.
162. Bals, R., D.J. Weiner, and J.M. Wilson, *The innate immune system in cystic fibrosis lung disease*. J Clin Invest, 1999. **103**(3): p. 303-7.
163. Lasky, L.A., *Selectins: interpreters of cell-specific carbohydrate information during inflammation*. Science, 1992. **258**(5084): p. 964-9.
164. Mahdavi, J., et al., *Helicobacter pylori SabA adhesin in persistent infection and chronic inflammation*. Science, 2002. **297**(5581): p. 573-8.
165. Davril, M., et al., *The sialylation of bronchial mucins secreted by patients suffering from cystic fibrosis or from chronic bronchitis is related to the severity of airway infection*. Glycobiology, 1999. **9**(3): p. 311-21.
166. Traving, C. and R. Schauer, *Structure, function and metabolism of sialic acids*. Cellular and molecular life sciences : CMLS, 1998. **54**(12): p. 1330-49.
167. Tonnaer, E.L., et al., *Involvement of glycosaminoglycans in the attachment of pneumococci to nasopharyngeal epithelial cells*. Microbes Infect, 2006. **8**(2): p. 316-22.
168. Bidossi, A., et al., *A functional genomics approach to establish the complement of carbohydrate transporters in Streptococcus pneumoniae*. PLoS One, 2012. **7**(3): p. e33320.
169. King, S.J., K.R. Hippe, and J.N. Weiser, *Deglycosylation of human glycoconjugates by the sequential activities of exoglycosidases expressed by Streptococcus pneumoniae*. Molecular microbiology, 2006. **59**(3): p. 961-74.

170. Trappetti, C., et al., *Sialic acid: a preventable signal for pneumococcal biofilm formation, colonization, and invasion of the host*. J Infect Dis, 2009. **199**(10): p. 1497-505.
171. Siegel, S.J., A.M. Roche, and J.N. Weiser, *Influenza Promotes Pneumococcal Growth during Coinfection by Providing Host Sialylated Substrates as a Nutrient Source*. Cell host & microbe, 2014. **16**(1): p. 55-67.
172. Marion, C., et al., *Sialic acid transport contributes to pneumococcal colonization*. Infect Immun, 2011. **79**(3): p. 1262-9.
173. Hatcher, B.L., J.Y. Hale, and D.E. Briles, *Free Sialic Acid Acts as a Signal That Promotes Streptococcus pneumoniae Invasion of Nasal Tissue and Nonhematogenous Invasion of the Central Nervous System*. Infect Immun, 2016. **84**(9): p. 2607-15.
174. Rogers, G.N. and J.C. Paulson, *Receptor determinants of human and animal influenza virus isolates: differences in receptor specificity of the H3 hemagglutinin based on species of origin*. Virology, 1983. **127**(2): p. 361-73.
175. Baum, L.G. and J.C. Paulson, *Sialyloligosaccharides of the respiratory epithelium in the selection of human influenza virus receptor specificity*. Acta Histochem Suppl, 1990. **40**: p. 35-8.
176. Wessels, M.R., et al., *Definition of a bacterial virulence factor: sialylation of the group B streptococcal capsule*. Proc Natl Acad Sci U S A, 1989. **86**(22): p. 8983-7.
177. Carlin, A.F., et al., *Molecular mimicry of host sialylated glycans allows a bacterial pathogen to engage neutrophil Siglec-9 and dampen the innate immune response*. Blood, 2009. **113**(14): p. 3333-6.
178. King, S.J., et al., *Phase variable desialylation of host proteins that bind to Streptococcus pneumoniae in vivo and protect the airway*. Molecular microbiology, 2004. **54**(1): p. 159-71.
179. Chang, Y.C., et al., *Leukocyte inflammatory responses provoked by pneumococcal sialidase*. mBio, 2012. **3**(1).
180. Grewal, P.K., et al., *The Ashwell receptor mitigates the lethal coagulopathy of sepsis*. Nat Med, 2008. **14**(6): p. 648-55.
181. Mahla, R.S., et al., *Sweeten PAMPs: Role of Sugar Complexed PAMPs in Innate Immunity and Vaccine Biology*. Front Immunol, 2013. **4**: p. 248.
182. Wood, D.M., et al., *Effect of hyperglycaemia on glucose concentration of human nasal secretions*. Clin Sci (Lond), 2004. **106**(5): p. 527-33.
183. Plumbridge, J. and E. Vimr, *Convergent pathways for utilization of the amino sugars N-acetylglucosamine, N-acetylmannosamine, and N-acetylneuraminic acid by Escherichia coli*. J Bacteriol, 1999. **181**(1): p. 47-54.
184. Vimr, E.R., et al., *Diversity of microbial sialic acid metabolism*. Microbiology and molecular biology reviews : MMBR, 2004. **68**(1): p. 132-53.
185. Vimr, E.R. and F.A. Troy, *Identification of an inducible catabolic system for sialic acids (nan) in Escherichia coli*. Journal of bacteriology, 1985. **164**(2): p. 845-53.
186. Hopkins, A.P., J.A. Hawkhead, and G.H. Thomas, *Transport and catabolism of the sialic acids N-glycolylneuraminic acid and 3-keto-3-deoxy-D-glycero-D-*

- galactonononic acid by Escherichia coli K-12*. FEMS Microbiol Lett, 2013. **347**(1): p. 14-22.
187. Almagro-Moreno, S. and E.F. Boyd, *Insights into the evolution of sialic acid catabolism among bacteria*. BMC evolutionary biology, 2009. **9**: p. 118.
  188. Tettelin, H., et al., *Complete genome sequence of a virulent isolate of Streptococcus pneumoniae*. Science, 2001. **293**(5529): p. 498-506.
  189. Gualdi, L., et al., *Regulation of neuraminidase expression in Streptococcus pneumoniae*. BMC microbiology, 2012. **12**: p. 200.
  190. Kostyukova, N.N., et al., *A study of pathogenic factors of Streptococcus pneumoniae strains causing meningitis*. FEMS Immunol Med Microbiol, 1995. **10**(2): p. 133-7.
  191. Watson, D.A. and D.M. Musher, *Interruption of capsule production in Streptococcus pneumonia serotype 3 by insertion of transposon Tn916*. Infect Immun, 1990. **58**(9): p. 3135-8.
  192. Melin, M., et al., *The capsular serotype of Streptococcus pneumoniae is more important than the genetic background for resistance to complement*. Infect Immun, 2010. **78**(12): p. 5262-70.
  193. Melin, M., et al., *Serotype-related variation in susceptibility to complement deposition and opsonophagocytosis among clinical isolates of Streptococcus pneumoniae*. Infect Immun, 2010. **78**(12): p. 5252-61.
  194. Hammerschmidt, S., *Adherence molecules of pathogenic pneumococci*. Current opinion in microbiology, 2006. **9**(1): p. 12-20.
  195. Weiser, J.N., et al., *Phase variation in pneumococcal opacity: relationship between colonial morphology and nasopharyngeal colonization*. Infect Immun, 1994. **62**(6): p. 2582-9.
  196. Weiser, J.N., *Phase variation in colony opacity by Streptococcus pneumoniae*. Microb Drug Resist, 1998. **4**(2): p. 129-35.
  197. Burnaugh, A.M., L.J. Frantz, and S.J. King, *Growth of Streptococcus pneumoniae on human glycoconjugates is dependent upon the sequential activity of bacterial exoglycosidases*. Journal of bacteriology, 2008. **190**(1): p. 221-30.
  198. Pettigrew, M.M., et al., *Variation in the presence of neuraminidase genes among Streptococcus pneumoniae isolates with identical sequence types*. Infection and immunity, 2006. **74**(6): p. 3360-5.
  199. Carvalho, S.M., et al., *CcpA ensures optimal metabolic fitness of Streptococcus pneumoniae*. PLoS One, 2011. **6**(10): p. e26707.
  200. Afzal, M., et al., *Sialic acid-mediated gene expression in Streptococcus pneumoniae and role of NanR as a transcriptional activator of the nan gene cluster*. Applied and environmental microbiology, 2015. **81**(9): p. 3121-31.
  201. Tong, H.H., et al., *Effect of neuraminidase on receptor-mediated adherence of Streptococcus pneumoniae to chinchilla tracheal epithelium*. Acta oto-laryngologica, 2002. **122**(4): p. 413-9.
  202. Camara, M., et al., *A neuraminidase from Streptococcus pneumoniae has the features of a surface protein*. Infect Immun, 1994. **62**(9): p. 3688-95.



203. Uchiyama, S., et al., *The surface-anchored NanA protein promotes pneumococcal brain endothelial cell invasion*. The Journal of experimental medicine, 2009. **206**(9): p. 1845-52.
204. Banerjee, A., et al., *Activation of brain endothelium by pneumococcal neuraminidase NanA promotes bacterial internalization*. Cell Microbiol, 2010. **12**(11): p. 1576-88.
205. Tong, H.H., et al., *Comparison of structural changes of cell surface carbohydrates in the eustachian tube epithelium of chinchillas infected with a Streptococcus pneumoniae neuraminidase-deficient mutant or its isogenic parent strain*. Microb Pathog, 2001. **31**(6): p. 309-17.
206. Shakhnovich, E.A., S.J. King, and J.N. Weiser, *Neuraminidase expressed by Streptococcus pneumoniae desialylates the lipopolysaccharide of Neisseria meningitidis and Haemophilus influenzae: a paradigm for interbacterial competition among pathogens of the human respiratory tract*. Infect Immun, 2002. **70**(12): p. 7161-4.
207. LeMessurier, K.S., A.D. Ogunniyi, and J.C. Paton, *Differential expression of key pneumococcal virulence genes in vivo*. Microbiology, 2006. **152**(Pt 2): p. 305-11.
208. Oggioni, M.R., et al., *Switch from planktonic to sessile life: a major event in pneumococcal pathogenesis*. Mol Microbiol, 2006. **61**(5): p. 1196-210.
209. Manco, S., et al., *Pneumococcal neuraminidases A and B both have essential roles during infection of the respiratory tract and sepsis*. Infection and immunity, 2006. **74**(7): p. 4014-20.
210. Orihuela, C.J., et al., *Tissue-specific contributions of pneumococcal virulence factors to pathogenesis*. The Journal of infectious diseases, 2004. **190**(9): p. 1661-9.
211. Blanchette, K.A., et al., *Neuraminidase A-Exposed Galactose Promotes Streptococcus pneumoniae Biofilm Formation during Colonization*. Infect Immun, 2016. **84**(10): p. 2922-32.
212. Berry, A.M. and J.C. Paton, *Additive attenuation of virulence of Streptococcus pneumoniae by mutation of the genes encoding pneumolysin and other putative pneumococcal virulence proteins*. Infect Immun, 2000. **68**(1): p. 133-40.
213. Chen, G.Y., et al., *Amelioration of sepsis by inhibiting sialidase-mediated disruption of the CD24-SiglecG interaction*. Nat Biotechnol, 2011. **29**(5): p. 428-35.
214. Stock, J.B., A.J. Ninfa, and A.M. Stock, *Protein phosphorylation and regulation of adaptive responses in bacteria*. Microbiological reviews, 1989. **53**(4): p. 450-90.
215. Throup, J.P., et al., *A genomic analysis of two-component signal transduction in Streptococcus pneumoniae*. Molecular microbiology, 2000. **35**(3): p. 566-76.
216. Guenzi, E., et al., *A two-component signal-transducing system is involved in competence and penicillin susceptibility in laboratory mutants of Streptococcus pneumoniae*. Molecular microbiology, 1994. **12**(3): p. 505-15.
217. Mascher, T., et al., *The CiaRH system of Streptococcus pneumoniae prevents lysis during stress induced by treatment with cell wall inhibitors and by mutations in pbp2x involved in beta-lactam resistance*. Journal of bacteriology, 2006. **188**(5): p. 1959-68.
218. Mascher, T., et al., *The Streptococcus pneumoniae cia regulon: CiaR target sites and transcription profile analysis*. Journal of bacteriology, 2003. **185**(1): p. 60-70.

219. Sebert, M.E., et al., *Microarray-based identification of htrA, a Streptococcus pneumoniae gene that is regulated by the CiaRH two-component system and contributes to nasopharyngeal colonization*. Infection and immunity, 2002. **70**(8): p. 4059-67.
220. Ibrahim, Y.M., et al., *Control of virulence by the two-component system CiaR/H is mediated via HtrA, a major virulence factor of Streptococcus pneumoniae*. Journal of bacteriology, 2004. **186**(16): p. 5258-66.
221. Giammarinaro, P., M. Sicard, and A.M. Gasc, *Genetic and physiological studies of the CiaH-CiaR two-component signal-transducing system involved in cefotaxime resistance and competence of Streptococcus pneumoniae*. Microbiology, 1999. **145** (Pt 8): p. 1859-69.
222. Peterson, S.N., et al., *Identification of competence pheromone responsive genes in Streptococcus pneumoniae by use of DNA microarrays*. Mol Microbiol, 2004. **51**(4): p. 1051-70.
223. Marx, P., et al., *Identification of genes for small non-coding RNAs that belong to the regulon of the two-component regulatory system CiaRH in Streptococcus*. BMC Genomics, 2010. **11**: p. 661.
224. Halfmann, A., et al., *Identification of the genes directly controlled by the response regulator CiaR in Streptococcus pneumoniae: five out of 15 promoters drive expression of small non-coding RNAs*. Molecular microbiology, 2007. **66**(1): p. 110-26.
225. Schnorpfel, A., et al., *Target evaluation of the non-coding csRNAs reveals a link of the two-component regulatory system CiaRH to competence control in Streptococcus pneumoniae R6*. Mol Microbiol, 2013. **89**(2): p. 334-49.
226. Tsui, H.C., et al., *Identification and characterization of noncoding small RNAs in Streptococcus pneumoniae serotype 2 strain D39*. J Bacteriol, 2010. **192**(1): p. 264-79.
227. Mann, B., et al., *Control of virulence by small RNAs in Streptococcus pneumoniae*. PLoS Pathog, 2012. **8**(7): p. e1002788.
228. Halfmann, A., et al., *Activity of the two-component regulatory system CiaRH in Streptococcus pneumoniae R6*. Journal of molecular microbiology and biotechnology, 2011. **20**(2): p. 96-104.
229. Pericone, C.D., et al., *Factors contributing to hydrogen peroxide resistance in Streptococcus pneumoniae include pyruvate oxidase (SpxB) and avoidance of the toxic effects of the fenton reaction*. Journal of bacteriology, 2003. **185**(23): p. 6815-25.
230. Marx, P., M. Meiers, and R. Bruckner, *Activity of the response regulator CiaR in mutants of Streptococcus pneumoniae R6 altered in acetyl phosphate production*. Frontiers in microbiology, 2014. **5**: p. 772.
231. Spellerberg, B., et al., *Pyruvate oxidase, as a determinant of virulence in Streptococcus pneumoniae*. Molecular microbiology, 1996. **19**(4): p. 803-13.
232. Pericone, C.D., et al., *Inhibitory and bactericidal effects of hydrogen peroxide production by Streptococcus pneumoniae on other inhabitants of the upper respiratory tract*. Infect Immun, 2000. **68**(7): p. 3990-7.

233. Avery, O.T. and H.J. Morgan, *The Occurrence of Peroxide in Cultures of Pneumococcus*. J Exp Med, 1924. **39**(2): p. 275-87.
234. Ibrahim, Y.M., et al., *Role of HtrA in the virulence and competence of Streptococcus pneumoniae*. Infection and immunity, 2004. **72**(6): p. 3584-91.
235. McCleary, W.R., J.B. Stock, and A.J. Ninfa, *Is acetyl phosphate a global signal in Escherichia coli?* J Bacteriol, 1993. **175**(10): p. 2793-8.
236. Feldman, C., et al., *The effects of pneumolysin and hydrogen peroxide, alone and in combination, on human ciliated epithelium in vitro*. Respir Med, 2002. **96**(8): p. 580-5.
237. Braun, J.S., et al., *Pneumococcal pneumolysin and H<sub>2</sub>O<sub>2</sub> mediate brain cell apoptosis during meningitis*. J Clin Invest, 2002. **109**(1): p. 19-27.
238. Rai, P., et al., *Streptococcus pneumoniae secretes hydrogen peroxide leading to DNA damage and apoptosis in lung cells*. Proc Natl Acad Sci U S A, 2015. **112**(26): p. E3421-30.
239. Loose, M., et al., *Pneumococcal hydrogen peroxide-induced stress signaling regulates inflammatory genes*. J Infect Dis, 2015. **211**(2): p. 306-16.
240. Echlin, H., et al., *Pyruvate Oxidase as a Critical Link between Metabolism and Capsule Biosynthesis in Streptococcus pneumoniae*. PLoS Pathog, 2016. **12**(10): p. e1005951.
241. Sebert, M.E., et al., *Pneumococcal HtrA protease mediates inhibition of competence by the CiaRH two-component signaling system*. Journal of bacteriology, 2005. **187**(12): p. 3969-79.
242. Spiess, C., A. Beil, and M. Ehrmann, *A temperature-dependent switch from chaperone to protease in a widely conserved heat shock protein*. Cell, 1999. **97**(3): p. 339-47.
243. Gee, J.B., et al., *Catalase-dependent peroxidative metabolism in the alveolar macrophage during phagocytosis*. The Journal of clinical investigation, 1970. **49**(6): p. 1280-7.
244. Rossi, F. and M. Zatti, *Biochemical aspects of phagocytosis in polymorphonuclear leucocytes. NADH and NADPH oxidation by the granules of resting and phagocytizing cells*. Experientia, 1964. **20**(1): p. 21-3.
245. Cassone, M., et al., *The HtrA protease from Streptococcus pneumoniae digests both denatured proteins and the competence-stimulating peptide*. J Biol Chem, 2012. **287**(46): p. 38449-59.
246. Boehm, M., et al., *Rapid paracellular transmigration of Campylobacter jejuni across polarized epithelial cells without affecting TER: role of proteolytic-active HtrA cleaving E-cadherin but not fibronectin*. Gut Pathog, 2012. **4**(1): p. 3.
247. Hoy, B., et al., *Helicobacter pylori HtrA is a new secreted virulence factor that cleaves E-cadherin to disrupt intercellular adhesion*. EMBO Rep, 2010. **11**(10): p. 798-804.
248. de Stoppelaar, S.F., et al., *Streptococcus pneumoniae serine protease HtrA, but not SFP or PrtA, is a major virulence factor in pneumonia*. PLoS One, 2013. **8**(11): p. e80062.

249. Libman, E., *A pneumococcus producing a peculiar form of hemolysis*. Proc. NY Pathol. Soc, 1905(5): p. 168.
250. Berry, A.M., et al., *Reduced virulence of a defined pneumolysin-negative mutant of Streptococcus pneumoniae*. Infect Immun, 1989. **57**(7): p. 2037-42.
251. Alouf, J.E., *Cholesterol-binding cytolytic protein toxins*. Int J Med Microbiol, 2000. **290**(4-5): p. 351-6.
252. Tilley, S.J., et al., *Structural basis of pore formation by the bacterial toxin pneumolysin*. Cell, 2005. **121**(2): p. 247-56.
253. Lim, J.E., et al., *Characterization of pneumolysin from Streptococcus pneumoniae, interacting with carbohydrate moiety and cholesterol as a component of cell membrane*. Biochem Biophys Res Commun, 2013. **430**(2): p. 659-63.
254. Shewell, L.K., et al., *The cholesterol-dependent cytolytins pneumolysin and streptolysin O require binding to red blood cell glycans for hemolytic activity*. Proc Natl Acad Sci U S A, 2014. **111**(49): p. E5312-20.
255. Spooncer, E., et al., *Isolation and characterization of polyfucosylated lactosaminoglycan from human granulocytes*. J Biol Chem, 1984. **259**(8): p. 4792-801.
256. Fukuda, M., et al., *Structure of sialylated fucosyl lactosaminoglycan isolated from human granulocytes*. J Biol Chem, 1984. **259**(17): p. 10925-35.
257. Johnson, M.K., *Cellular location of pneumolysin*. FEMS Microbiol Lett, 1977. **2**: p. 243-245.
258. Walker, J.A., et al., *Molecular cloning, characterization, and complete nucleotide sequence of the gene for pneumolysin, the sulfhydryl-activated toxin of Streptococcus pneumoniae*. Infect Immun, 1987. **55**(5): p. 1184-9.
259. Price, K.E. and A. Camilli, *Pneumolysin localizes to the cell wall of Streptococcus pneumoniae*. J Bacteriol, 2009. **191**(7): p. 2163-8.
260. Balachandran, P., et al., *The autolytic enzyme LytA of Streptococcus pneumoniae is not responsible for releasing pneumolysin*. J Bacteriol, 2001. **183**(10): p. 3108-16.
261. Benton, K.A., J.C. Paton, and D.E. Briles, *The hemolytic and complement-activating properties of pneumolysin do not contribute individually to virulence in a pneumococcal bacteremia model*. Microb Pathog, 1997. **23**(4): p. 201-9.
262. Price, K.E., N.G. Greene, and A. Camilli, *Export requirements of pneumolysin in Streptococcus pneumoniae*. J Bacteriol, 2012. **194**(14): p. 3651-60.
263. Rubins, J.B., et al., *Pneumolysin in pneumococcal adherence and colonization*. Microb Pathog, 1998. **25**(6): p. 337-42.
264. Houldsworth, S., P.W. Andrew, and T.J. Mitchell, *Pneumolysin stimulates production of tumor necrosis factor alpha and interleukin-1 beta by human mononuclear phagocytes*. Infect Immun, 1994. **62**(4): p. 1501-3.
265. Ferrante, A., B. Rowan-Kelly, and J.C. Paton, *Inhibition of in vitro human lymphocyte response by the pneumococcal toxin pneumolysin*. Infect Immun, 1984. **46**(2): p. 585-9.

266. Littmann, M., et al., *Streptococcus pneumoniae evades human dendritic cell surveillance by pneumolysin expression*. EMBO Mol Med, 2009. **1**(4): p. 211-22.
267. Dessing, M.C., et al., *Role of Toll-like receptors 2 and 4 in pulmonary inflammation and injury induced by pneumolysin in mice*. PLoS One, 2009. **4**(11): p. e7993.
268. Canvin, J.R., et al., *The role of pneumolysin and autolysin in the pathology of pneumonia and septicemia in mice infected with a type 2 pneumococcus*. J Infect Dis, 1995. **172**(1): p. 119-23.
269. Zafar, M.A., et al., *Host-to-Host Transmission of Streptococcus pneumoniae Is Driven by Its Inflammatory Toxin, Pneumolysin*. Cell Host Microbe, 2017. **21**(1): p. 73-83.
270. Pittet, L.A., et al., *Influenza virus infection decreases tracheal mucociliary velocity and clearance of Streptococcus pneumoniae*. Am J Respir Cell Mol Biol, 2010. **42**(4): p. 450-60.
271. McCullers, J.A. and K.C. Bartmess, *Role of neuraminidase in lethal synergism between influenza virus and Streptococcus pneumoniae*. J Infect Dis, 2003. **187**(6): p. 1000-9.
272. Plotkowski, M.C., et al., *Adherence of type I Streptococcus pneumoniae to tracheal epithelium of mice infected with influenza A/PR8 virus*. Am Rev Respir Dis, 1986. **134**(5): p. 1040-4.
273. Ghoneim, H.E., P.G. Thomas, and J.A. McCullers, *Depletion of alveolar macrophages during influenza infection facilitates bacterial superinfections*. J Immunol, 2013. **191**(3): p. 1250-9.
274. Sun, K. and D.W. Metzger, *Inhibition of pulmonary antibacterial defense by interferon-gamma during recovery from influenza infection*. Nat Med, 2008. **14**(5): p. 558-64.
275. Sun, K. and D.W. Metzger, *Influenza infection suppresses NADPH oxidase-dependent phagocytic bacterial clearance and enhances susceptibility to secondary methicillin-resistant Staphylococcus aureus infection*. J Immunol, 2014. **192**(7): p. 3301-7.
276. Rynda-Appl, A., K.M. Robinson, and J.F. Alcorn, *Influenza and Bacterial Superinfection: Illuminating the Immunologic Mechanisms of Disease*. Infect Immun, 2015. **83**(10): p. 3764-70.
277. Lacks, S. and R.D. Hotchkiss, *A study of the genetic material determining an enzyme in Pneumococcus*. Biochimica et biophysica acta, 1960. **39**: p. 508-18.
278. Mellroth, P., et al., *LytA, major autolysin of Streptococcus pneumoniae, requires access to nascent peptidoglycan*. J Biol Chem, 2012. **287**(14): p. 11018-29.
279. Karreman, C., *Fusion PCR, a one-step variant of the "megaprimer" method of mutagenesis*. Biotechniques, 1998. **24**(5): p. 736, 740, 742.
280. Pruss, B.M. and A.J. Wolfe, *Regulation of acetyl phosphate synthesis and degradation, and the control of flagellar expression in Escherichia coli*. Molecular microbiology, 1994. **12**(6): p. 973-84.
281. Pan, X., Y. Yang, and J.R. Zhang, *Molecular basis of host specificity in human pathogenic bacteria*. Emerging microbes & infections, 2014. **3**(3): p. e23.

282. Hedlund, M., et al., *N-glycolylneuraminic acid deficiency in mice: implications for human biology and evolution*. Mol Cell Biol, 2007. **27**(12): p. 4340-6.
283. Bardor, M., et al., *Mechanism of uptake and incorporation of the non-human sialic acid N-glycolylneuraminic acid into human cells*. J Biol Chem, 2005. **280**(6): p. 4228-37.
284. Oetke, C., et al., *Evidence for efficient uptake and incorporation of sialic acid by eukaryotic cells*. Eur J Biochem, 2001. **268**(16): p. 4553-61.
285. Chiavolini, D., G. Pozzi, and S. Ricci, *Animal models of Streptococcus pneumoniae disease*. Clinical microbiology reviews, 2008. **21**(4): p. 666-85.
286. Mahajan, V.S., et al., *Striking Immune Phenotypes in Gene-Targeted Mice Are Driven by a Copy-Number Variant Originating from a Commercially Available C57BL/6 Strain*. Cell Rep, 2016. **15**(9): p. 1901-9.
287. Feldman, C., et al., *The interaction of Streptococcus pneumoniae with intact human respiratory mucosa in vitro*. The European respiratory journal, 1992. **5**(5): p. 576-83.
288. Makatsori, E., et al., *Screening of N-acetylneuraminic acids in serum and tissue specimens of mouse C57Bl with Lewis' lung cancer by high-performance liquid chromatography*. Journal of chromatography. B, Biomedical sciences and applications, 1998. **712**(1-2): p. 23-9.
289. Minami, A., et al., *Catalytic preference of Salmonella typhimurium LT2 sialidase for N-acetylneuraminic acid residues over N-glycolylneuraminic acid residues*. FEBS open bio, 2013. **3**: p. 231-6.
290. Parker, R.B., J.E. McCombs, and J.J. Kohler, *Sialidase specificity determined by chemoselective modification of complex sialylated glycans*. ACS chemical biology, 2012. **7**(9): p. 1509-14.
291. Corfield, A.P., et al., *The release of N-acetyl- and N-glycoloyl-neuraminic acid from soluble complex carbohydrates and erythrocytes by bacterial, viral and mammalian sialidases*. The Biochemical journal, 1981. **197**(2): p. 293-9.
292. Marra, A., et al., *Differential fluorescence induction analysis of Streptococcus pneumoniae identifies genes involved in pathogenesis*. Infection and immunity, 2002. **70**(3): p. 1422-33.
293. Paixao, L., et al., *Transcriptional and metabolic effects of glucose on Streptococcus pneumoniae sugar metabolism*. Frontiers in microbiology, 2015. **6**: p. 1041.
294. Kingston, G.W., P.T. Phang, and M.J. Leathley, *Increased incidence of nosocomial pneumonia in mechanically ventilated patients with subclinical aspiration*. American journal of surgery, 1991. **161**(5): p. 589-92.
295. Han, B.G., et al., *Markers of glycemic control in the mouse: comparisons of 6-h- and overnight-fasted blood glucoses to Hb A1c*. American journal of physiology. Endocrinology and metabolism, 2008. **295**(4): p. E981-6.
296. Syk, A., et al., *Emergence of hypervirulent mutants resistant to early clearance during systemic serotype 1 pneumococcal infection in mice and humans*. J Infect Dis, 2014. **210**(1): p. 4-13.
297. Spellerberg, B., et al., *Pneumococcal cell wall activates NF-kappa B in human monocytes: aspects distinct from endotoxin*. Microbial pathogenesis, 1996. **20**(5): p. 309-17.

298. Zychlinsky, A. and P.J. Sansonetti, *Apoptosis as a proinflammatory event: what can we learn from bacteria-induced cell death?* Trends Microbiol, 1997. **5**(5): p. 201-4.
299. Shrestha, S., et al., *Identifying the interaction between influenza and pneumococcal pneumonia using incidence data.* Sci Transl Med, 2013. **5**(191): p. 191ra84.
300. McCullers, J.A. and J.E. Rehg, *Lethal synergism between influenza virus and Streptococcus pneumoniae: characterization of a mouse model and the role of platelet-activating factor receptor.* J Infect Dis, 2002. **186**(3): p. 341-50.
301. McCullers, J.A., *Effect of antiviral treatment on the outcome of secondary bacterial pneumonia after influenza.* J Infect Dis, 2004. **190**(3): p. 519-26.
302. Shahangian, A., et al., *Type I IFNs mediate development of postinfluenza bacterial pneumonia in mice.* J Clin Invest, 2009. **119**(7): p. 1910-20.
303. McNamee, L.A. and A.G. Harmsen, *Both influenza-induced neutrophil dysfunction and neutrophil-independent mechanisms contribute to increased susceptibility to a secondary Streptococcus pneumoniae infection.* Infect Immun, 2006. **74**(12): p. 6707-21.
304. Robinson, K.M., et al., *Influenza A virus exacerbates Staphylococcus aureus pneumonia in mice by attenuating antimicrobial peptide production.* J Infect Dis, 2014. **209**(6): p. 865-75.
305. Chromek, M., et al., *The antimicrobial peptide cathelicidin protects the urinary tract against invasive bacterial infection.* Nat Med, 2006. **12**(6): p. 636-41.
306. Buffinton, G.D., et al., *Oxidative stress in lungs of mice infected with influenza A virus.* Free Radic Res Commun, 1992. **16**(2): p. 99-110.
307. Wong, S.M., et al., *Genome-wide fitness profiling reveals adaptations required by Haemophilus in coinfection with influenza A virus in the murine lung.* Proc Natl Acad Sci U S A, 2013. **110**(38): p. 15413-8.
308. Hendriksen, W.T., et al., *Regulation of gene expression in Streptococcus pneumoniae by response regulator 09 is strain dependent.* J Bacteriol, 2007. **189**(4): p. 1382-9.
309. Sun, K., et al., *Interleukin-12 promotes gamma interferon-dependent neutrophil recruitment in the lung and improves protection against respiratory Streptococcus pneumoniae infection.* Infect Immun, 2007. **75**(3): p. 1196-202.
310. Haraguchi, S., et al., *Interleukin 12 deficiency associated with recurrent infections.* Proc Natl Acad Sci U S A, 1998. **95**(22): p. 13125-9.
311. Bernatoniene, J., et al., *Induction of CC and CXC chemokines in human antigen-presenting dendritic cells by the pneumococcal proteins pneumolysin and CbpA, and the role played by toll-like receptor 4, NF-kappaB, and mitogen-activated protein kinases.* J Infect Dis, 2008. **198**(12): p. 1823-33.
312. Shoma, S., et al., *Critical involvement of pneumolysin in production of interleukin-1alpha and caspase-1-dependent cytokines in infection with Streptococcus pneumoniae in vitro: a novel function of pneumolysin in caspase-1 activation.* Infect Immun, 2008. **76**(4): p. 1547-57.
313. Srivastava, A., et al., *The apoptotic response to pneumolysin is Toll-like receptor 4 dependent and protects against pneumococcal disease.* Infect Immun, 2005. **73**(10): p. 6479-87.

- 314. Ratner, A.J., et al., *Epithelial cells are sensitive detectors of bacterial pore-forming toxins*. J Biol Chem, 2006. **281**(18): p. 12994-8.
- 315. Kuri, T., et al., *Influenza A virus-mediated priming enhances cytokine secretion by human dendritic cells infected with Streptococcus pneumoniae*. Cell Microbiol, 2013. **15**(8): p. 1385-400.
- 316. Heinz, S., et al., *Species-specific regulation of Toll-like receptor 3 genes in men and mice*. J Biol Chem, 2003. **278**(24): p. 21502-9.
- 317. Tanabe, M., et al., *Mechanism of up-regulation of human Toll-like receptor 3 secondary to infection of measles virus-attenuated strains*. Biochem Biophys Res Commun, 2003. **311**(1): p. 39-48.
- 318. Bai, W., et al., *TLR3 regulates mycobacterial RNA-induced IL-10 production through the PI3K/AKT signaling pathway*. Cell Signal, 2014. **26**(5): p. 942-50.
- 319. Hovden, A.O., et al., *The bacterial preparation OK432 induces IL-12p70 secretion in human dendritic cells in a TLR3 dependent manner*. PLoS One, 2012. **7**(2): p. e31217.
- 320. Kawashima, T., et al., *Double-stranded RNA of intestinal commensal but not pathogenic bacteria triggers production of protective interferon-beta*. Immunity, 2013. **38**(6): p. 1187-97.